



**OPPA European Social Fund
Prague & EU: We invest in your future.**

AE4M33RZN, Fuzzy logic: Fuzzy relations

Radomír Černoč

radomir.cernoch@fel.cvut.cz

26/11/2012

Faculty of Electrical Engineering, CTU in Prague

Plan of the lecture

Properties of fuzzy sets

- Fuzzy implication and fuzzy properties

- Fuzzy set inclusion and crisp predicates

Intermission: Probabilistic vs. fuzzy

Binary fuzzy relations

- Quick revision of crisp relations

- Fuzzyfication of crisp relations

- Projection and cylindrical extension

- Composition of fuzzy relations

- Properties of fuzzy relations

- Properties of fuzzy composition

Extensions

Biblography

Organizational:

- Next week, there will be a short test (max 5 points).
- This week we are having the last **theoretical lecture**.

Fuzzy implication

We already know *fuzzy negation* \neg_{\circ} , *fuzzy conjunction* \wedge_{\circ} and *fuzzy disjunction* \vee_{\circ} . What about other operators?

Fuzzy implication

We already know *fuzzy negation* $\overset{\circ}{\neg}$, *fuzzy conjunction* $\overset{\circ}{\wedge}$ and *fuzzy disjunction* $\overset{\circ}{\vee}$. What about other operators?

Definition

Fuzzy implication is any function

$$\overset{\circ}{\Rightarrow} : [0, 1]^2 \rightarrow [0, 1] \quad (1)$$

which overlaps with the boolean implication on $x, y \in \{0, 1\}$:

$$(x \overset{\circ}{\Rightarrow} y) = (x \Rightarrow y) . \quad (2)$$

Residue implication

Despite the lack of a uniform definition of fuzzy implication, there is a useful class of implications:

Defintion



The *R-implication* (residuum, „*reziduovaná implikace*“) is a function obtained from a fuzzy T-norm:

$$\alpha \overset{R}{\underset{\circ}{\Rightarrow}} \beta = \sup\{\gamma \mid \alpha \underset{\circ}{\wedge} \gamma \leq \beta\} \quad (\text{RI})$$

R-implication: Examples (1)

Standard implication (Gödel) is derived from (RI) using the standard conjunction \wedge_S :



$$\alpha \xrightarrow[S]{R} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ \beta & \text{otherwise} \end{cases} \quad (3)$$



R-implication: Examples (2)

Lukasiewicz implication is derived from (RI) using the Lukasiewicz conjunction \wedge_L :



$$\alpha \xrightarrow[\text{L}]{\text{R}} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ 1 - \alpha + \beta & \text{otherwise} \end{cases} \quad (4)$$



R-implication: Examples (3)

Algebraic implication (Gougen, Gaines) is derived from (RI) using the algebraic conjunction \wedge_A :

$$\alpha \xrightarrow[A]{R} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ \frac{\beta}{\alpha} & \text{otherwise} \end{cases} \quad (5)$$



R-implication: Properties

Theorem 209.

Let \triangleleft_{\circ} be a continuous fuzzy conjunction. Then R-implication satisfies:

$$\alpha \xrightarrow[\circ]{R} \beta = \mathbf{1} \text{ iff } \alpha \leq \beta \quad (I1)$$

$$\mathbf{1} \xrightarrow[\circ]{R} \beta = \beta \quad (I2)$$

$$\alpha \xrightarrow[\circ]{R} \beta \text{ is not increasing in } \alpha \text{ and not decreasing in } \beta \quad (I3)$$

R-implication: Properties

Proof of theorem 209.

Let's denote $\{\gamma \mid \alpha \underset{\circ}{\wedge} \gamma \leq \beta\} = \gamma$.

- Proving (I3) uses monotonicity: Increasing α can only shrink γ and increasing β can only enlarge γ .
- Proving (I2) is easy: $1 \xrightarrow{\underset{\circ}{R}} \beta = \sup\{\gamma \mid 1 \underset{\circ}{\wedge} \gamma \leq \beta\}$. From definition of $\underset{\circ}{\wedge}$, we write $1 \xrightarrow{\underset{\circ}{R}} \beta = \sup\{\gamma \mid \gamma \leq \beta\} = \beta$.

Proof of theorem 209 (contd.).

- For (I1) one needs to check 2 cases:
 - If $\alpha \leq \beta$, then $1 \in \gamma$, because $\alpha \wedge_{\circ} 1 = \alpha \leq \beta$ and therefore the condition $\alpha \wedge_{\circ} \gamma \leq \beta$ is true for all possible values of γ .
 - If $\alpha > \beta$, then $1 \notin \gamma$, because $\alpha \wedge_{\circ} 1 = \alpha > \beta$ and therefore the condition $\alpha \wedge_{\circ} \gamma \leq \beta$ is false for $\gamma = 1$.

S-implication

Defintion

The *S-implication* is a function obtained from a fuzzy disjunction $\overset{\circ}{\vee}$:

$$\alpha \underset{\circ}{\overset{\text{S}}{\Rightarrow}} \beta = \underset{\text{S}}{\neg} \alpha \overset{\circ}{\vee} \beta \quad (\text{SI})$$

S-implication

Definition

The *S-implication* is a function obtained from a fuzzy disjunction $\overset{\circ}{\vee}$:

$$\alpha \overset{\text{S}}{\underset{\circ}{\Rightarrow}} \beta = \underset{\text{S}}{\neg} \alpha \overset{\circ}{\vee} \beta \quad (\text{SI})$$

Example

Kleene-Dienes implication from $\overset{\text{S}}{\underset{\circ}{\vee}}$

$$\alpha \overset{\text{S}}{\underset{\circ}{\Rightarrow}} \beta = \max(1 - \alpha, \beta) \quad (6)$$

Generalized fuzzy inclusion

Generalized fuzzy inclusion

Previously, we used the logical negation \neg to define the set complement, the conjunction \wedge to define the set intersection, etc.

Can we use the implication \Rightarrow to define the the fuzzy inclusion?

Generalized fuzzy inclusion

Previously, we used the logical negation \neg to define the set complement, the conjunction \wedge to define the set intersection, etc.

Can we use the implication \Rightarrow to define the the fuzzy inclusion?

Definition

The *generalized fuzzy inclusion* $\overset{\circ}{\subseteq}$ is a function that assigns a degree to the the inclusion of set $A \in \mathbb{F}(\Delta)$ in set $B \in \mathbb{F}(\Delta)$:

$$A \overset{\circ}{\subseteq} B = \inf\{A(\mathbf{x}) \overset{\circ}{\Rightarrow} B(\mathbf{x}) \mid \mathbf{x} \in \Delta\} \quad (7)$$

Generalized fuzzy inclusion: Example

Fuzzy inclusion (non-generalized)

Definition

The *fuzzy inclusion* \subseteq is a predicate (assigns a true/false value) which hold for two fuzzy sets $A, B \in \mathbb{F}(\Delta)$ iff

$$\mu_A(\mathbf{x}) \leq \mu_B(\mathbf{x}) \text{ for all } \mathbf{x} \in \Delta. \quad (8)$$

Fuzzy inclusion (non-generalized)

In vertical representation, the definition has a straightforward equivalent:

$$\mu_A \leq \mu_B \quad (9)$$

Fuzzy inclusion (non-generalized)

In vertical representation, the definition has a straightforward equivalent:

$$\mu_A \leq \mu_B \quad (9)$$

In horizontal representation, there is a theorem:

Theorem 219.

Let $A, B \in \mathbb{F}(\Delta)$ if and only if

$$R_A(\alpha) \subseteq R_B(\alpha) \text{ for all } \alpha \in [0, 1] . \quad (10)$$

Fuzzy inclusion (non-generalized)

Proof of theorem 219.

- \Rightarrow Assume $A \subseteq B$ and $\mathbf{x} \in R_A(\alpha)$ for some value α . If $\alpha \leq A(\mathbf{x})$, then $A(\mathbf{x}) \leq B(\mathbf{x})$ (from the definition of $A \subseteq B$) and therefore $\mathbf{x} \in R_B(\alpha)$ and $R_A(\alpha) \subseteq R_B(\alpha)$.
- \Leftarrow Assume $R_A(\alpha) \subseteq R_B(\alpha)$. Firstly recall the horizontal-vertical translation formula: $\mu_A(\mathbf{x}) = \sup\{\alpha \in [0, 1] \mid \mathbf{x} \in R_A(\alpha)\}$. Since $\{\alpha \mid \mathbf{x} \in R_A(\mathbf{x})\} \subseteq \{\alpha \mid \mathbf{x} \in R_B(\mathbf{x})\}$, the inequality $A(\mathbf{x}) \leq \sup\{\alpha \mid \mathbf{x} \in R_B(\mathbf{x})\} \leq B(\mathbf{x})$ holds.

Cutworthiness

We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

Cutworthiness

Let P be a predicate (returns true/false) over fuzzy sets. P is called *cutworthy* („řezově dědičná vlastnost“) if the implication holds:

$$P(A_1, \dots, A_n) \Rightarrow P(R_{A_1}(\alpha), \dots, R_{A_n}(\alpha)) \text{ for all } \alpha \in [0, 1] \quad (11)$$

We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

Cutworthiness

Let P be a predicate (returns true/false) over fuzzy sets. P is called *cutworthy* („řezově dědičná vlastnost“) if the implication holds:

$$P(A_1, \dots, A_n) \Rightarrow P(R_{A_1}(\alpha), \dots, R_{A_n}(\alpha)) \text{ for all } \alpha \in [0, 1] \quad (11)$$

There is a related notion: We define P as *cut-consistent* („řezově konzistentní“) using the same definition, but replacing \Rightarrow with \Leftrightarrow .

Cutworthiness: Examples

- The theorem 219 can be stated as: “Set inclusion is cut-consistent.”

Brain teasers

- *Strong normality* of A is defined as $A(x) = 1$ for some $x \in \Delta$.
????
- *Being crisp* is
????


Cutworthiness: Examples

- The theorem 219 can be stated as: “Set inclusion is cut-consistent.”

Brain teasers


- *Strong normality* of A is defined as $A(\mathbf{x}) = 1$ for some $\mathbf{x} \in \Delta$.
Strong normality is **cut-consistent**: A is strongly-normal iff every its cut is non-empty iff every cut strongly normal.
- *Being crisp* is **cutworthy, but not cut-consistent**: Every cut is crisp by definition, therefore cutworthiness. But even **non-crisp sets** have crisp cuts, therefore the property is not not cut-consistent.

Google: “fuzzy”



Sources: M. Taylor's Weblog, M. Taylor's Weblog, Eddie's Trick Shop.

Google: “probability”



Sources: Life123, WhatWeKnowSoFar, Probability Problems.

Fuzzy vs. probability

- *Vagueness vs. uncertainty.*

Fuzzy vs. probability

- *Vagueness vs. uncertainty.*
- Fuzzy logic is *functional*.

Definition

A *binary crisp relation* R from X onto Y is a subset of the cartesian product $X \times Y$:

$$R \in \mathbb{P}(X \times Y) \quad (12)$$

Definition

A *binary crisp relation* R from X onto Y is a subset of the cartesian product $X \times Y$:

$$R \in \mathbb{P}(X \times Y) \quad (12)$$

Definition

The *inverse relation* R^{-1} to R is a relation from Y to X s.t.

$$R^{-1} = \{(y, x) \in Y \times X \mid (x, y) \in R\} \quad (13)$$

Crisp relations: Inverse

Definition

Let X, Y, Z be sets. Then the *compound* of relations $R \subseteq X \times Y, S \subseteq Y \times Z$ is the relation

$$R \circ S = \{(x, z) \in X \times Z \mid (x, y) \in R \text{ and } (y, z) \in S \text{ for some } y\} \quad (14)$$

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property

using logical connectives

using set axioms

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$
transitive	$(x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$	$R \circ R \subseteq R$

Crisp relations: Properties

The *equality* relation on Δ is $E = \{(x, x) \mid x \in \Delta\}$.

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$
transitive	$(x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$	$R \circ R \subseteq R$
partial order	reflexive, transitive and anti-symmetric	
equivalence	reflexive, transitive and symmetric	

Definition

A *binary fuzzy relation* R from X onto Y is a fuzzy subset on the universe $X \times Y$.

$$R \in \mathbb{F}(X \times Y) \quad (15)$$

Definition

The *fuzzy inverse* relation $R^{-1} \in \mathbb{F}(Y \times X)$ to $R \in \mathbb{F}(X \times Y)$, s.t.

$$R(\mathbf{y}, \mathbf{x}) = R^{-1}(\mathbf{x}, \mathbf{y}) \quad (16)$$

Defintion

Let $R \in \mathbb{F}(X \times Y)$ be a fuzzy binary relation. The *first* and second projection of R is

$$R^{(1)}(\mathbf{x}) = \bigvee_{\mathbf{y} \in Y}^S R(\mathbf{x}, \mathbf{y}) \quad (17)$$

$$R^{(2)}(\mathbf{y}) = \bigvee_{\mathbf{x} \in X}^S R(\mathbf{x}, \mathbf{y}) \quad (18)$$

Projection: Example

R	y_1	y_2	y_3	y_4	y_5	y_6	$R^{(1)}(x)$
x_1	0.1	0.2	0.4	0.8	1	0.8	?
x_2	0.2	0.4	0.8	1	0.8	0.6	?
x_3	0.4	0.8	1	0.8	0.4	0.2	?
$R^{(2)}(y)$?	?	?	?	?	?	

Projection: Example

R	y_1	y_2	y_3	y_4	y_5	y_6	$R^{(1)}(x)$
x_1	0.1	0.2	0.4	0.8	1	0.8	1
x_2	0.2	0.4	0.8	1	0.8	0.6	1
x_3	0.4	0.8	1	0.8	0.4	0.2	1
$R^{(2)}(y)$	0.4	0.8	1	0.8	0.4	0.2	

Sometimes there is a *total projection* defined as

$R^{(T)} = \bigvee_{x \in X} \bigvee_{y \in Y} R(x, y)$. But we already know this notion as $\text{Height}(R)$.

Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

Definition

Let $A \in \mathbb{F}(X)$ and $B \in \mathbb{F}(Y)$ be fuzzy sets. The *cylindrical extension* („cylindrické rozšíření“, „kartézský součin fuzzy množin“) is defined as

$$A \times B(x, y) = A(x) \underset{S}{\wedge} B(y) \quad (19)$$

Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

Definition

Let $A \in \mathbb{F}(X)$ and $B \in \mathbb{F}(Y)$ be fuzzy sets. The *cylindrical extension* („cylindrické rozšíření“, „kartézský součin fuzzy množin“) is defined as

$$A \times B(x, y) = A(x) \underset{S}{\wedge} B(y) \quad (19)$$

Brain teaser

Why can't there be a relation Q bigger than $A \times B$, whose projections are $Q^{(1)} = A$ and $Q^{(2)} = B$?

Cylindrical extension: Drawing

$$A(x) = \begin{cases} x - 1 & x \in [1, 2] \\ 3 - x & x \in [2, 3] \\ 0 & \text{otherwise} \end{cases}$$

$$B(x) = \begin{cases} x - 3 & x \in [3, 4] \\ 5 - x & x \in [4, 5] \\ 0 & \text{otherwise} \end{cases}$$

Composition of fuzzy relations

Definition

Let X, Y, Z be crisp sets. $R \in \mathbb{F}(X \times Y)$, $S \in \mathbb{F}(Y \times Z)$ and \wedge some fuzzy conjunction. Then the \circ -composition („ \circ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge S(y, z) \quad (20)$$

Composition of fuzzy relations

Definition

Let X, Y, Z be crisp sets. $R \in \mathbb{F}(X \times Y)$, $S \in \mathbb{F}(Y \times Z)$ and \wedge some fuzzy conjunction. Then the \circ -composition („ \circ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge S(y, z) \quad (20)$$

1. For infinite domains, \bigvee^S is computed using the **sup** instead of **max**.

Composition of fuzzy relations

Definition

Let X, Y, Z be crisp sets. $R \in \mathbb{F}(X \times Y)$, $S \in \mathbb{F}(Y \times Z)$ and \wedge some fuzzy conjunction. Then the \circ -composition („ \circ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge S(y, z) \quad (20)$$

1. For infinite domains, \bigvee^S is computed using the \sup instead of \max .
2. Instead of the “for some y ” in *crisp relations*, the disjunction “finds such a y ” that maximizes the conjunction.

Example of a fuzzy relation

$$R(x, y) = \begin{cases} x + y & x, y \in [0, \frac{1}{2}] \\ 0 & \text{otherwise} \end{cases}$$

$$S(x, y) = \begin{cases} x \cdot y & x, y \in [0, 1] \\ 0 & \text{otherwise} \end{cases}$$

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property

using set axioms

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using set axioms
reflexive	$E \subseteq R$

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
\circ -anti-symmetric	$R \cap R^{-1} \subseteq E$

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
\circ -anti-symmetric	$R \cap_{\circ} R^{-1} \subseteq E$
\circ -transitive	$R \circ_{\circ} R \subseteq R$

Properties of fuzzy relations

Then the relation $R \subseteq \Delta \times \Delta$ is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
\circ -anti-symmetric	$R \cap_{\circ} R^{-1} \subseteq E$
\circ -transitive	$R \circ_{\circ} R \subseteq R$
\circ -partial order	reflexive, \circ -transitive and \circ -anti-symmetric
\circ -equivalence	reflexive, \circ -transitive and \circ -symmetric

Properties in a finite domain

If the universe Δ is finite, the relation can be written as a matrix. Their properties are reflected in the relation's matrix:

- **Reflexivity:** Cells on the main diagonal ?.
- **Symmetry:** Cells symmetric over the main diagonal ?.
- **Anti-symmetry:** Cells symmetric over the main diagonal ?.
 - For *S*- and *A*-anti-symmetry, ?.
 - For *L*-anti-symmetry, ?.
- **Transitivity:** More difficult (see example on the next slide).

Properties in a finite domain

If the universe Δ is finite, the relation can be written as a matrix. Their properties are reflected in the relation's matrix:

- **Reflexivity:** Cells on the main diagonal are 1.
- **Symmetry:** Cells symmetric over the main diagonal are equal.
- **Anti-symmetry:** Cells symmetric over the main diagonal have conjunction equal to zero.
 - For S- and A-anti-symmetry, one of the elements must be zero.
 - For L-anti-symmetry, their sum must be less or equal to 1.
- **Transitivity:** More difficult (see example on the next slide).

Example on fuzzy relation properties

Let $\Delta = \{A, B, C, D\}$ and $R \in \mathbb{F}(\Delta \times \Delta)$.

R	A	B	C	D
A		0.5		0.1
B			0.2	
C				
D		0.2		

Fill the missing cells in the table to make R

- S-equivalence
- A-equivalence

Theorem 264.

Let R , S and T be relations (defined over sets that “make sense”) The following equations hold:

Theorem 264.

Let R , S and T be relations (defined over sets that “make sense”) The following equations hold:

$$R \circ E = R, \quad E \circ R = R \quad (21)$$

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1} \quad (22)$$

$$R \circ (S \circ T) = (R \circ S) \circ T \quad (23)$$

$$\left(R \bigcap^S S \right) \circ T = (R \circ T) \circ (S \circ T) \quad (24)$$

$$R \circ \left(S \bigcap^S T \right) = (R \circ S) \circ (R \circ T) \quad (25)$$

Theorem 264.

Let R , S and T be relations (defined over sets that “make sense”) The following equations hold:

$$R \circ E = R, \quad E \circ R = R \quad (21)$$

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1} \quad (22)$$

$$R \circ (S \circ T) = (R \circ S) \circ T \quad (23)$$

$$\left(R \bigcap^S S \right) \circ T = (R \circ T) \circ (S \circ T) \quad (24)$$

$$R \circ \left(S \bigcap^S T \right) = (R \circ S) \circ (R \circ T) \quad (25)$$

(21) describes the *identity element*, (22) the *inverse of composition*, (23) is the *asociativity*, (24) and (25) the *right-* and *left-distributivity*.

Proof of 264.

Proving (21) and (22) is trivial.

$$"R \circ (S \circ T)"(x, w) = \bigvee_y^s R(x, y) \wedge "S \circ T"(y, w) \quad (26)$$

$$= \bigvee_y^s R(x, y) \wedge \left(\bigvee_z^s S(y, z) \wedge T(z, w) \right) \quad (27)$$

$$= \bigvee_y^s \bigvee_z^s R(x, y) \wedge S(y, z) \wedge T(z, w) \quad (28)$$

$$= \bigvee_z^s \bigvee_y^s R(x, y) \wedge S(y, z) \wedge T(z, w) \quad (29)$$

Proof of 264 (contd.).

$$= \bigvee_z^s \bigvee_y^s R(x, y) \underset{\circ}{\wedge} S(y, z) \underset{\circ}{\wedge} T(z, w) \quad (30)$$

$$= \bigvee_z^s \left(\bigvee_y^s R(x, y) \underset{\circ}{\wedge} S(y, z) \right) \underset{\circ}{\wedge} T(z, w) \quad (31)$$

$$= \bigvee_z^s "R \underset{\circ}{S}"(x, z) \underset{\circ}{\wedge} T(z, w) \quad (32)$$

$$= "R \underset{\circ}{S} \underset{\circ}{T}"(x, w) \quad (33)$$

Proof of (24) and (25) is similar (uses the distributivity law), only shorter. See [Navara and Olšák, 2001] for details.

Extensions: Sometimes it is useful to consider...

- ...a ε -*reflective* relation

$$R(\mathbf{x}, \mathbf{x}) \geq \varepsilon \quad (34)$$

Extensions: Sometimes it is useful to consider...

- ...a ε -*reflective* relation

$$R(\mathbf{x}, \mathbf{x}) \geq \varepsilon \quad (34)$$

- ...a *weakly reflexive* relation

$$R(\mathbf{x}, \mathbf{y}) \leq R(\mathbf{x}, \mathbf{x}) \text{ and } R(\mathbf{y}, \mathbf{x}) \leq R(\mathbf{x}, \mathbf{x}) \text{ for all } \mathbf{x}, \mathbf{y} \quad (35)$$

Extensions: Sometimes it is useful to consider...

- ...a ε -*reflective* relation

$$R(x, x) \geq \varepsilon \quad (34)$$

- ...a *weakly reflexive* relation

$$R(x, y) \leq R(x, x) \text{ and } R(y, x) \leq R(x, x) \text{ for all } x, y \quad (35)$$

- Relation is 1-reflective **iff** reflexive.
- If a relation is reflexive, **then** it is weakly reflexive.

Extensions: Sometimes it is useful to consider...

- ...a *non-involutive negation* by refusing (N2)

$$\neg \neg \alpha \neq \alpha$$

and adopting a weaker axiom

$$\neg \neg 0 = 1 \text{ and } \neg \neg 1 = 0 \quad (\text{N0})$$

Example

Gödel negation

$$\neg_G \alpha = \begin{cases} 1 & \alpha = 0 \\ 0 & \text{otherwise} \end{cases} \quad (36)$$



Navara, M. and Olšák, P. (2001).
Základy fuzzy množin.
Nakladatelství ČVUT.



**OPPA European Social Fund
Prague & EU: We invest in your future.**
