



**OPPA European Social Fund
Prague & EU: We invest in your future.**

► The Representation Theorem for Essential Matrices

Let SVD of E be UDV^T

Theorem

A 3×3 matrix E is an essential matrix iff $D \simeq \text{diag}(1, 1, 0)$.



Proof.

$$\tilde{E} = [e_2, e_1] R$$

1. Part I: General properties of antisymmetric 3×3 matrices

2. Part II (direct):

If E is essential then it has two equal singular values and the third is zero.

3. Part III (converse):

Let $A = \hat{U}D\hat{V}^T$ s.t. $D = \text{diag}(1, 1, 0)$ then $A = [\hat{u}_3]_{\times} R$, where R is orthogonal, \hat{u}_3 is the

3rd column of \hat{U} , and $R = \hat{U}W\hat{V}^T$, where $W = \begin{bmatrix} 0 & \alpha & 0 \\ -\alpha & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. $|\alpha| = 1$

□


Proof, Part I: More Properties of Antisymmetric 3×3 Matrices

Given vector \mathbf{b} , let there be matrices \mathbf{D} , \mathbf{W} , \mathbf{V}

$$\mathbf{D} = \|\mathbf{b}\| \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \mathbf{W} = \begin{bmatrix} 0 & \alpha & 0 \\ -\alpha & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \mathbf{V} = \left[\mathbf{a}, \mathbf{c}, \frac{\mathbf{b}}{\|\mathbf{b}\|} \right] \quad (11)$$

such that

1. $|\alpha| = 1$
2. $\|\mathbf{a}\| = \|\mathbf{c}\| = 1$
3. $\mathbf{a}, \mathbf{c}, \mathbf{b}$ mutually orthogonal: $\mathbf{V}^T \mathbf{V} = \mathbf{I}$
4. $\det \mathbf{V} = 1$



b is given, a, c not

note that

- $\mathbf{W}^T \mathbf{W} = \mathbf{I}$; \mathbf{W} is a rotation by 90°
- if $\alpha \mapsto -\alpha$ then $\mathbf{W} \mapsto \mathbf{W}^T$
- \mathbf{a}, \mathbf{c} are determined up to a rotation φ about \mathbf{b} , $\hat{\mathbf{V}} = \mathbf{T}_\varphi \mathbf{V}$, $\mathbf{T}_\varphi \mathbf{b} = \mathbf{b}$

Theorem (A)

Let \mathbf{V} , \mathbf{D} , \mathbf{W} , \mathbf{T}_φ be defined as above. Then $\hat{\mathbf{U}} \mathbf{D} \hat{\mathbf{V}}^T$ is an SVD of $[\mathbf{b}]_{\times}$ iff $\hat{\mathbf{U}} = \mathbf{T}_\varphi \mathbf{V} \mathbf{W}^T$, $\hat{\mathbf{V}} = \mathbf{T}_\varphi \mathbf{V}$ for some φ .

$$\mathbf{T}_\varphi^T \hat{\mathbf{V}} = \mathbf{V} \rightarrow \hat{\mathbf{U}} = \underbrace{\mathbf{T}_\varphi^T \mathbf{T}_\varphi^T}_{\mathbf{I}} \hat{\mathbf{V}} \mathbf{W}^T$$

It follows $\hat{\mathbf{U}} = \hat{\mathbf{V}} \mathbf{W}^T$ for any φ and $\hat{\mathbf{U}} \mathbf{D} \hat{\mathbf{V}}^T = \hat{\mathbf{V}} \mathbf{W}^T \mathbf{D} \hat{\mathbf{V}}^T = \hat{\mathbf{U}} \mathbf{D} \mathbf{W}^T \hat{\mathbf{U}}$

Proof of Theorem A.

1. Converse ($\hat{\mathbf{U}}, \hat{\mathbf{V}}, \mathbf{D}, \mathbf{V}, \mathbf{W}, \mathbf{T}_\varphi$ as defined $\Rightarrow \hat{\mathbf{U}}\mathbf{D}\hat{\mathbf{V}}^\top$ is an SVD of $[\mathbf{b}]_x$):

a. $\hat{\mathbf{U}}\mathbf{D}\hat{\mathbf{V}}^\top = \underbrace{\mathbf{T}_\varphi \mathbf{V}\mathbf{W}^\top}_{\hat{\mathbf{U}}} \mathbf{D} \underbrace{\mathbf{V}^\top \mathbf{T}_\varphi^\top}_{\hat{\mathbf{V}}^\top}$ is indeed an SVD of some matrix for any φ .

b. what matrix?

$$\mathbf{T}_\varphi \mathbf{V}\mathbf{W}^\top \mathbf{D}\mathbf{V}^\top \mathbf{T}_\varphi^\top = \mathbf{T}_\varphi \|\mathbf{b}\| (\mathbf{c}\mathbf{a}^\top - \mathbf{a}\mathbf{c}^\top) \mathbf{T}_\varphi^\top = \|\mathbf{b}\| \mathbf{T}_\varphi [\mathbf{a} \times \mathbf{c}]_x \mathbf{T}_\varphi^\top = \mathbf{T}_\varphi [\mathbf{b}]_x \mathbf{T}_\varphi^\top = [\mathbf{T}_\varphi \mathbf{b}]_x = [\mathbf{b}]_x \quad (12)$$

$\begin{bmatrix} 1 & 0 \\ 0 & \vdots \end{bmatrix} \mathbf{V}^\top \rightarrow \begin{bmatrix} \vdots & \vdots & \vdots \\ 0 & 0 & 0 \end{bmatrix}$

hence it is an SVD of $[\mathbf{b}]_x$ but also of $[\mathbf{T}_\varphi \mathbf{b}]_x$ for any φ

2. Direct: For every φ we go backward in (12) and obtain an SVD.

$$(\mathbf{c}\mathbf{a}^\top - \mathbf{a}\mathbf{c}^\top) \tilde{\mathbf{b}} \stackrel{?}{=} \mathbf{0} \quad \mathbf{v} = [\mathbf{a}, \mathbf{c}, \frac{\mathbf{b}}{\|\mathbf{b}\|}] \quad \mathbf{v}\mathbf{v}^\top = \mathbf{I} \quad \square$$

$$\underbrace{\mathbf{c}\mathbf{a}^\top}_{\mathbf{0}} - \underbrace{\mathbf{a}\mathbf{c}^\top}_{\mathbf{1}} = -\mathbf{a}$$

We are proving (from Slide 78):

Part II

If \mathbf{E} is essential then it has two equal singular values and the third is zero.

- The \mathbf{E} is essential, hence $\mathbf{E} \simeq [\mathbf{t}]_{\times} \mathbf{R}$ $t = -t_2$
- Let $\hat{\mathbf{U}}\mathbf{D}\hat{\mathbf{V}}^{\top}$ be the SVD of $[\mathbf{t}]_{\times}$. Then, by Theorem A, $\underbrace{\hat{\mathbf{U}}}_{\text{orthogonal}} \mathbf{D} \underbrace{\hat{\mathbf{V}}^{\top} \mathbf{R}}_{\text{orthogonal}}$ is an SVD of \mathbf{E} with singular values $\mathbf{D} = \underbrace{\|\mathbf{t}\|}_{\wedge} \text{diag}(1, 1, 0)$.

Part III

Let $\mathbf{A} = \hat{\mathbf{U}}\mathbf{D}\hat{\mathbf{V}}^{\top}$ s.t. $\mathbf{D} = \text{diag}(1, 1, 0)$ then $\mathbf{A} = [\hat{\mathbf{u}}_3]_{\times} \mathbf{R}$, where \mathbf{R} is orthogonal.

$$\hat{\mathbf{U}}\mathbf{D} \underbrace{\hat{\mathbf{V}}^{\top}}_{\text{choice: } \mathbf{W}\hat{\mathbf{U}}^{\top} \mathbf{R}} = \underbrace{\hat{\mathbf{U}}\mathbf{D}\mathbf{W}\hat{\mathbf{U}}^{\top}}_{\text{antisymmetric by Theorem A}} \mathbf{R} = [\hat{\mathbf{u}}_3]_{\times} \mathbf{R}$$

where $[\hat{\mathbf{u}}_3]_{\times}$ is obtained by inspection and we have defined $\hat{\mathbf{V}}$ s.t. $\mathbf{R} = \hat{\mathbf{U}}\mathbf{W}\hat{\mathbf{V}}^{\top}$

► Essential Matrix Decomposition

Essential matrix captures relative camera position

[Longuet-Higgins 1981]

$$\mathbf{E} = [-\mathbf{t}_{21}]_{\times} \mathbf{R}_{21} = [\mathbf{R}_2 \mathbf{b}]_{\times} \mathbf{R}_{21} = \mathbf{R}_{21} [\mathbf{R}_1 \mathbf{b}]_{\times}$$

- rank $\mathbf{E} = 2$ since rank $[\mathbf{t}_{21}]_{\times} = 2$
- Let $\mathbf{E} = \mathbf{U} \mathbf{D} \mathbf{V}^{\top}$ be the SVD of \mathbf{E} s.t. $\mathbf{D} = \text{diag}(1, 1, 0)$. Then [H&Z, sec. 9.6]
 - in case $\det \mathbf{U} < 0$ transform it to $-\mathbf{U}$, do the same for \mathbf{V}
 - compute

$$\mathbf{R}_{21} = \mathbf{U} \begin{bmatrix} 0 & \alpha & 0 \\ -\alpha & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{V}^{\top}, \quad \mathbf{t}_{21} = -\mathbf{U} \begin{bmatrix} 0 \\ 0 \\ \beta \end{bmatrix}, \quad |\alpha| = 1, \quad \beta \neq 0 \quad (13)$$

$$[U, D, V] = \text{svd}(E)$$

$$\text{if } \det(U) < 0 \\ U = -U; V = -V$$

$$|\alpha| = 1, \quad \beta \neq 0$$

end

$$\mathbf{R}_{21} = \mathbf{U} \mathbf{W} \mathbf{V}^{\top}$$

$$\mathbf{t}_{21} = -\mu(:, 3)$$

despite non-uniqueness of SVD


$$\mathbf{R}_{21}' = \mathbf{U} \mathbf{W}^{\top} \mathbf{V}^{\top}$$

$$\mathbf{t}_{21}' = +\mu(:, 3)$$

Notes

- the result for \mathbf{R}_{21} is unique up to $\alpha = \pm 1$
- change of sign in \mathbf{W} rotates the solution by 180° about \mathbf{t}
 $\mathbf{R}_1 = \mathbf{U} \mathbf{W} \mathbf{V}^{\top}$, $\mathbf{R}_2 = \mathbf{U} \mathbf{W}^{\top} \mathbf{V}^{\top} \Rightarrow \mathbf{T} = \mathbf{R}_2 \mathbf{R}_1^{\top} = \dots = \mathbf{U} \text{diag}(-1, -1, 1) \mathbf{U}^{\top}$ which is a rotation by 180° about $\mathbf{u}_3 = \mathbf{t}_{21}$
- \mathbf{t}_{21} recoverable up to scale β and direction sign β
- 4 solution sets for 4 sign combinations of α, β see next for geometric interpretation

► Four Solutions to Essential Matrix Decomposition



- chirality constraint: all 3D points are in front of both cameras
- this singles-out the upper left case

[H&Z, Sec. 9.6.3]

► 7-Point Algorithm for Estimating Fundamental Matrix

Problem: Given a set $\{(x_i, y_i)\}_{i=1}^k$ of $k = 7$ correspondences, estimate f. m. \mathbf{F} .

$$\underline{\mathbf{y}}_i^\top \mathbf{F} \underline{\mathbf{x}}_i = 0, \quad i = 1, \dots, k, \quad \text{known: } \underline{\mathbf{x}}_i = (x_{i1}, x_{i2}, 1), \quad \underline{\mathbf{y}}_i = (y_{i1}, y_{i2}, 1)$$

terminology: correspondence = truth, later: match = algorithm's result; hypothesised corresp.

Solution:

$$\mathbf{D} = \begin{matrix} & \overset{1 \times 6}{x_{11}y_{11}} & & \overset{1 \times 3}{x_{11}} & & & & & \overset{1 \times 0}{1} \\ \begin{bmatrix} x_{11}y_{11} & x_{11}y_{12} & x_{11} & x_{12}y_{11} & x_{12}y_{12} & x_{12} & y_{11} & y_{12} & 1 \\ x_{21}y_{21} & x_{21}y_{22} & x_{21} & x_{22}y_{21} & x_{22}y_{22} & x_{22} & y_{21} & y_{22} & 1 \\ \vdots & & & & & & & & \vdots \\ x_{k1}y_{k1} & x_{k1}y_{k2} & x_{k1} & x_{k2}y_{k1} & x_{k2}y_{k2} & x_{k2} & y_{k1} & y_{k2} & 1 \end{bmatrix}, & \mathbf{D} \in \mathbb{R}^{k,9}
 \end{matrix}$$

$$\overset{7 \times 9}{\mathbf{D}} \mathbf{f} = \mathbf{0}, \quad \mathbf{f} = [f_{11} \quad f_{21} \quad f_{31} \quad \dots \quad f_{33}]^\top, \quad \mathbf{f} \in \mathbb{R}^9,$$

- for $k = 7$ we have a rank-deficient system, the null-space of \mathbf{D} is 2-dimensional
- but we know that $\det \mathbf{F} = 0$
- 7-point algorithm:

1. find a basis of the null space of \mathbf{D} : $\mathbf{F}_1, \mathbf{F}_2$

by SVD or QR factorization

2. get up to 3 real solutions for α from

$$\det(\alpha \mathbf{F}_1 + (1 - \alpha) \mathbf{F}_2) = 0 \quad \text{cubic equation in } \alpha$$

3. get up to 3 fundamental matrices $\mathbf{F} = \alpha_i \mathbf{F}_1 + (1 - \alpha_i) \mathbf{F}_2$



- the result may depend on image transformations
- normalization improves conditioning
- this gives a good starting point for the full algorithm

Slide 91

Slide 110

► Degenerate Configurations for Fundamental Matrix Estimation


When is \mathbf{F} not uniquely determined from any number of correspondences? [H&Z, Sec. 11.9]

1. camera centers coincide $C_1 = C_2$



- epipolar geometry is not defined
- images are related by homography \mathbf{H}
- we do get an \mathbf{F} from the 7-point algorithm but it is of the form of $\mathbf{F} = \mathbf{S}\mathbf{H}$, with \mathbf{S} antisymmetric

$$y \approx Hx$$



$$l \approx \underline{s} \times \mathbf{H}\underline{x} \quad \text{arbitrary } s$$

$$y \in l: 0 = \underline{y}^T (\underline{s} \times \mathbf{H}\underline{x}) = \underline{y}^T \underbrace{[\underline{s}]_{\times} \mathbf{H}}_{\mathbf{S}\mathbf{H}} \underline{x}$$
$$y^T \mathbf{F} \underline{x} = 0$$

2. all 3D points lie in a plane

- images related by homography
- again, \mathbf{F} is not unique, $\mathbf{F} = \mathbf{S}\mathbf{H}$, where \mathbf{S} is as above

note essential matrix estimation can deal with planes, Slide 87

3. both camera centers and all 3D points lie on a ruled quadric

hyperboloid of one sheet, cones, cylinders, two planes

- there are 3 solutions for \mathbf{F}


notes

- a complete treatment with additional degenerate configurations in [H&Z, sec. 22.2]
- stronger epipolar constraint can reject some configurations
- we assume correct correspondences, dealing with mismatches need not be a part of the 7-point algorithm

→ Slide 112

A Note on Oriented Epipolar Constraint

- a tighter epipolar constraint preserves orientations
- requires all points and cameras be on the same side of the plane at infinity



$$\mathbf{e}_2 \times \mathbf{m}_2 \stackrel{\pm}{\sim} \mathbf{F} \mathbf{m}_1$$

notation: $\mathbf{m} \stackrel{\pm}{\sim} \mathbf{n}$ means $\mathbf{m} = \lambda \mathbf{n}$, $\lambda > 0$

- note that the constraint is not invariant to the change of either sign of \mathbf{m}_i
- all 7 correspondence in 7-point alg. must have the same sign
- this may help reject some wrong matches, see Slide 112
- an even more tight constraint: scene points in front of both cameras

see later

[Chum et al. 2004]

expensive

this is called chirality constraint

► Five-Point Algorithm for Relative Camera Orientation

Problem: Given $\{\underline{\mathbf{m}}_i, \underline{\mathbf{m}}'_i\}_{i=1}^5$ corresponding image points and calibration matrix \mathbf{K} , recover the camera motion \mathbf{R}, \mathbf{t} .

Obs:

1. \mathbf{R} – 3DOF, \mathbf{t} – we can recover 2DOF only, in total 5 DOF \rightarrow we need 3 constraints on \mathbf{E}
2. real $\mathbf{F} \in \mathbb{R}^{3,3}$ is a fundamental matrix iff $\det \mathbf{F} = 0$
3. fundamental matrix is essential iff its two non-zero eigenvalues are equal

This gives an equation system:

$$\begin{array}{ll} \underline{\mathbf{v}}_i^\top \mathbf{E} \underline{\mathbf{v}}'_i = 0 & \text{5 linear constraints } (\underline{\mathbf{v}} = \mathbf{K}^{-1} \underline{\mathbf{m}}) \\ \text{equal singular values } \downarrow \det \mathbf{E} = 0 & \text{1 cubic constraint} \\ \mathbf{E} \mathbf{E}^\top \mathbf{E} - \frac{1}{2} \text{tr}(\mathbf{E} \mathbf{E}^\top) \mathbf{E} = 0 & \text{9 cubic constraints, 2 independent} \end{array} \quad \left. \vphantom{\begin{array}{l} \\ \\ \end{array}} \right\} 8+1$$

1. estimate \mathbf{E} by SVD from $\underline{\mathbf{v}}_i^\top \mathbf{E} \underline{\mathbf{v}}'_i = 0$ by the null-space method, this gives $\mathbf{E} = x \mathbf{E}_1 + y \mathbf{E}_2 + z \mathbf{E}_3 + \mathbf{E}_4$
2. at most 10 (complex) solutions for x, y, z from the cubic constraints

- when all 3D points lie on a plane: at most 2 solutions (twisted-pair)

can be disambiguated in 3 views
or by chirality constraint (Slide 83) unless all 3D points are closer to one camera

- 6-point problem for unknown f [Kukelova et al. BMVC 2008]
- resources at http://cmp.felk.cvut.cz/minimal/5_pt_relative.php



**OPPA European Social Fund
Prague & EU: We invest in your future.**
