

**GRAPHICAL MARKOV MODELS (WS2017)**  
**5. SEMINAR**

**Assignment 1.** Consider the task of finding the most probable sequence of (hidden) states for a (Hidden) Markov model on a chain.

**a)** Show that the Dynamic Programming approach applied for this task can be interpreted as an equivalent transformation (re-parametrisation) of the model.

**b)** Show that the transformed functions (potentials) encode an explicit description of *all* optimisers of the problem

**Assignment 2.** Consider a GRF for binary valued labellings  $x: V \rightarrow \{0, 1\}$  of a graph  $(V, E)$  given by

$$p(x) = \frac{1}{Z} \exp \left[ \sum_{ij \in E} u_{ij}(x_i, x_j) + \sum_{i \in V} u_i(x_i) \right].$$

Show that it is always possible to find an equivalent transformation (re-parametrisation)

$$u_{ij} \rightarrow \tilde{u}_{ij}, \quad u_i \rightarrow \tilde{u}_i$$

such that the new pairwise functions  $\tilde{u}_{ij}$  have the form

$$\tilde{u}_{ij}(x_i, x_j) = \alpha_{ij} |x_i - x_j|$$

with some real numbers  $\alpha_{ij} \in \mathbb{R}$ .

**Assignment 3.** Transform the *Travelling Salesman Problem* into a (min, +)-problem.

**Assignment 4.** Let  $K$  be a completely ordered finite set. We assume w.l.o.g. that  $K = \{1, 2, \dots, m\}$ . For a function  $u: K \rightarrow \mathbb{R}$  define its discrete “derivative” by  $Du(k) = u(k+1) - u(k)$ .

**a)** Let  $u$  be a function  $u: K^2 \rightarrow \mathbb{R}$  and denote by  $D_1$  and  $D_2$  the discrete derivatives w.r.t. its first and second argument. Prove the following equality

$$D_1 D_2 u(k_1, k_2) = u(k_1 + 1, k_2 + 1) + u(k_1, k_2) - u(k_1 + 1, k_2) - u(k_1, k_2 + 1).$$

Conclude that all mixed derivatives  $D_1 D_2 u(k_1, k_2)$  of a submodular functions are negative.

**b)** Prove that the condition established in a) is not only necessary but also sufficient for a function to be submodular.

*Hint:* Start from the observation that the following equality holds for a function of one variable

$$u(k+l) - u(k) = \sum_{i=k}^{k+l-1} Du(i)$$

and generalise it for functions of two variables.

**c)** Prove that any function  $u: K^2 \rightarrow \mathbb{R}$  can be represented as a sum of a submodular and a supermodular function.

*Hint:* Consider the mixed derivative  $D_1 D_2 u(k_1, k_2)$ , decompose it into its negative and positive part and “integrate” them back separately.