

Quiz

Michal Štolba

stolba@agents.fel.cvut.cz



PUI (Planning in Artificial Intelligence)

Heuristic h dominates h' when

1. $h \leq h'$ for all states
2. $h \geq h'$ for all states
3. $h \neq h'$ for all states

Heuristic h dominates h' when

1. $h \leq h'$ for all states
2. $h \geq h'$ for all states
3. $h \neq h'$ for all states

What is the most commonly used relaxation of a STRIPS action $a = \langle \text{pre}(a), \text{add}(a), \text{del}(a) \rangle$?

1. $a^+ = \langle \emptyset, \text{add}(a), \text{del}(a) \rangle$
2. $a^+ = \langle \text{pre}(a), \emptyset, \text{del}(a) \rangle$
3. $a^+ = \langle \text{pre}(a), \text{add}(a), \emptyset \rangle$

What is the most commonly used relaxation of a STRIPS action $a = \langle \text{pre}(a), \text{add}(a), \text{del}(a) \rangle$?

1. $a^+ = \langle \emptyset, \text{add}(a), \text{del}(a) \rangle$
2. $a^+ = \langle \text{pre}(a), \emptyset, \text{del}(a) \rangle$
3. $a^+ = \langle \text{pre}(a), \text{add}(a), \emptyset \rangle$

How can a multi-valued variable $V \in \{v_1, \dots, v_k\}$ be relaxed?

How can a multi-valued variable $V \in \{v_1, \dots, v_k\}$ be relaxed?

Answer: Accumulating semantics: $V \subseteq \{v_1, \dots, v_k\}$.

What is not true?

1. Relaxations of plans are relaxed plans.
2. Relaxations are no harder to solve than the original task.
3. Optimal relaxed plans may be longer than optimal plans for original tasks.

What is not true?

1. Relaxations of plans are relaxed plans.
2. Relaxations are no harder to solve than the original task.
3. Optimal relaxed plans **may be** longer than the optimal plans for original tasks.

Answer: Optimal relaxed plans **are always shorter (or equal)** than the optimal plans for original tasks.

What heuristic is defined by

$$h(p) = \min_{a \in \text{add}(p)} h(a)$$

$$h(a) = \text{cost}(a) + \max_{p \in \text{pre}(a)} h(p)$$

1. h_{max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

What heuristic is defined by

$$h(p) = \min_{a \in \text{add}(p)} h(a)$$

$$h(a) = \text{cost}(a) + \max_{p \in \text{pre}(a)} h(p)$$

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

What heuristic is defined by

$$h(p) = \min_{a \in \text{add}(p)} h(a)$$

$$h(a) = \text{cost}(a) + \sum_{p \in \text{pre}(a)} h(p)$$

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

What heuristic is defined by

$$h(p) = \min_{a \in \text{add}(p)} h(a)$$

$$h(a) = \text{cost}(a) + \sum_{p \in \text{pre}(a)} h(p)$$

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

For h_{add} is not true:

1. Assumes independence of facts.
2. All facts are assumed to be achieved by the cheapest action.
3. Is admissible.

For h_{add} is not true:

1. Assumes independence of facts.
2. All facts are assumed to be achieved by the cheapest action.
3. Is admissible.

Answer: h_{add} is not admissible.

Why?

For h_{add} is not true:

1. Assumes independence of facts.
2. All facts are assumed to be achieved by the cheapest action.
3. Is admissible.

Answer: h_{add} is not admissible.

Why?

Which heuristic is admissible?

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

Which heuristic is admissible?

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

Why?

Why not h_{STRIPS} ?

Why not h_{FF} ?

Which heuristic is admissible?

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

Why?

Why not h_{STRIPS} ?

Why not h_{FF} ?

Which heuristic is admissible?

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

Why?

Why not h_{STRIPS} ?

Why not h_{FF} ?

Which heuristic is admissible?

1. h_{\max}
2. h_{add}
3. h_{STRIPS}
4. h_{FF}

Why?

Why not h_{STRIPS} ?

Why not h_{FF} ?