


Logical reasoning and programming, lab session X

(December 2, 2019)

- X.1** We have a language that contains only one binary predicate symbol \in and we have an interpretation $\mathcal{M} = (D, i)$ such that $D = \{a, b, c, d\}$ and $i(\in)$ is given by the following diagram:



Meaning that $x \in y$ iff there is an arrow from x to y . Decide whether the following formulae are valid in \mathcal{M} :

- (a) $\exists X \forall Y (\neg(Y \in X))$,
- (b) $\exists X \forall Y (Y \in X)$,
- (c) $\exists X \forall Y (Y \in X \leftrightarrow Y \in Y)$,
- (d) $\exists X \forall Y (Y \in X \leftrightarrow \neg(Y \in Y))$.

- X.2** Show that the following formulae are valid and provide counter-examples for the opposite implications:

- (a) $\forall X p(X) \vee \forall X q(X) \rightarrow \forall X (p(X) \vee q(X))$,
- (b) $\exists X (p(X) \wedge q(X)) \rightarrow \exists X p(X) \wedge \exists X q(X)$,
- (c) $\exists X \forall Y p(X, Y) \rightarrow \forall Y \exists X p(X, Y)$,
- (d) $\forall X p(X) \rightarrow \exists X p(X)$.

- X.3** Show that the “exists unique” quantifier $\exists!$ does not commute with \exists , \forall , nor $\exists!$.

- X.4** Decide whether for any formula φ holds:

- (a) $\varphi \equiv \forall \varphi$,
- (b) $\varphi \equiv \exists \varphi$,
- (c) $\models \varphi$ iff $\models \forall \varphi$,
- (d) $\models \varphi$ iff $\models \exists \varphi$,

where $\forall \varphi$ ($\exists \varphi$) is the universal (existential) closure of φ . If not, does at least one implication hold?

- X.5** Show that for any set of formulae Γ and a formula φ holds if $\Gamma \models \varphi$, then $\forall \Gamma \models \varphi$, where $\forall \Gamma = \{\forall \psi : \psi \in \Gamma\}$. Does the opposite direction hold?

- X.6** Find a set of formulae Γ and a formula φ such that $\Gamma \models \varphi$ and $\Gamma \not\models \neg \varphi$.

- X.7** Produce equivalent formulae in prenex form:

- (a) $\forall X (p(X) \rightarrow \forall Y (q(X, Y) \rightarrow \neg \forall Z r(Y, Z)))$,

- (b) $\exists X p(X, Y) \rightarrow (q(X) \rightarrow \neg \forall Z p(X, Z))$,
- (c) $\exists X p(X, Y) \rightarrow (q(X) \rightarrow \neg \exists Z p(X, Z))$,
- (d) $p(X, Y) \rightarrow \exists Y (q(Y) \rightarrow (\exists X q(X) \rightarrow r(Y)))$,
- (e) $\forall Y p(Y) \rightarrow (\forall X q(X) \rightarrow r(Z))$.

X.8 In **X.7** you could obtain in some cases various prefixes; the order of quantifiers can be different. Are all these variants correct?

X.9 Can we produce a formula equivalent to **X.7e** with just one quantifier?

X.10 Produce Skolemized formulae equisatisfiable with those in **X.7**. Try to produce as simple as possible Skolem functions.

X.11 Skolemize the following formula

$$\forall X (p(a) \vee \exists Y (q(Y) \wedge \forall Z (p(Y, Z) \vee \exists U q(X, Y)))) \vee \exists W q(a, W).$$

Why is it possible in this particular case to do that without producing an equivalent formula in prenex form?

X.12 Unify the following pairs of formulae:

- (a) $\{p(X, Y) \doteq p(Y, f(Z))\}$,
- (b) $\{p(a, Y, f(Y)) \doteq p(Z, Z, U)\}$,
- (c) $\{p(X, g(X)) \doteq p(Y, Y)\}$,
- (d) $\{p(X, g(X), Y) \doteq p(Z, U, g(U))\}$,
- (e) $\{p(g(X), Y) \doteq p(Y, Y), p(Y, Y) \doteq p(U, f(W))\}$.

X.13 What is the size of the maximal term that is produced when you try to unify

$$\{f(g(X_1, X_1), g(X_2, X_2), \dots, g(X_{n-1}, X_{n-1})) \doteq f(X_2, X_3, \dots, X_n)\}.$$