

# Probabilistic classification

Tomáš Svoboda and Matěj Hoffmann  
thanks to, Daniel Novák and Filip Železný

Vision for Robots and Autonomous Systems, Center for Machine Perception  
Department of Cybernetics  
Faculty of Electrical Engineering, Czech Technical University in Prague


May 6, 2020

# (Re-)introduction uncertainty/probability

- ▶ Markov Decision Processes - uncertainty about outcome of **actions**
- ▶ Now: uncertainty may be also associated with **states**
  - ▶ Different states may have different **prior probabilities**
  - ▶ The states  $s \in S$  may not be directly observable
  - ▶ They need to be inferred from **features**  $x \in X$
- ▶ This is addressed by the rules of probability (*such as Bayes theorem*) and leads on to
  - ▶ Bayesian classification
  - ▶ Bayesian decision making

## Probability example: Picking fruits

- ▶ red box: 2 apples, 6 oranges
- ▶ blue box: 3 apples, 1 orange




- ▶ Scenario: Pick a box (say red box in 40% cases), then pick a fruit at random
- ▶ (Frequent) questions:
  - ▶ What is the overall probability that the selection procedure will pick an apple?
  - ▶ Given that we have chosen an orange, what is the probability that the box we chose was the blue one?

Example from Chapter 1.2 [1]

# Picking fruits. What is the probability that ...?

- ▶ red box: 2 apples, 6 oranges
- ▶ blue box: 3 apples, 1 orange



Scenario: Pick a box (say red box in 40% cases), then pick a fruit at random.  
What is the overall probability that the selection procedure will pick an apple?

- A: 11/20
- B: 6/8
- C: 1/2
- D: Different value.

4 / 24

## Notes

Example serves for probability recap (sum, product rules, conditional probabilities, Bayes)

Random variables:

- Identity of the box  $B$ , two possible values  $r, b$
- Identity of the fruit  $F$ , two possible values  $a, o$

Info about picking a box  $P(B = r) = 0.4$  and  $P(B = b) = 0.6$ .

Conditional probabilities, given box selected:  $P(o|r) = 3/4$ ,  $P(a|r) = 1/4$ ,  $P(o|b) = 1/4$ ,  $P(a|b) = 3/4$ .

Answering questions:


- $P(F = a) = P(a|r)P(r) + P(a|b)P(b) = 11/20$
- $P(B = b|F = o) = P(b|o)$

$$P(b|o) = \frac{P(o|b)P(b)}{P(o)} = \frac{P(o|b)P(b)}{P(o|b)P(b) + P(o|r)P(r)} = 1/3$$

$P(B)$  prior probability - before we observe the fruit;  $P(B|F)$  - aposteriori probability - after we observe the fruit.

# Picking fruits. What is the probability that ...?

- ▶ red box: 2 apples, 6 oranges
- ▶ blue box: 3 apples, 1 orange



Scenario: Pick a box (say red box in 40% cases), then pick a fruit at random. What is the overall probability that the selection procedure will pick an apple?

- A:  $11/20$
- B:  $6/8$
- C:  $1/2$
- D: Different value.

What is the probability that the selection procedure will pick an apple?

## Notes

Example serves for probability recap (sum, product rules, conditional probabilities, Bayes)

Random variables:

- Identity of the box  $B$ , two possible values  $r, b$
- Identity of the fruit  $F$ , two possible values  $a, o$

Info about picking a box  $P(B = r) = 0.4$  and  $P(B = b) = 0.6$ .

Conditional probabilities, given box selected:  $P(o|r) = 3/4$ ,  $P(a|r) = 1/4$ ,  $P(o|b) = 1/4$ ,  $P(a|b) = 3/4$ .

Answering questions:


- $P(F = a) = P(a|r)P(r) + P(a|b)P(b) = 11/20$
- $P(B = b|F = o) = P(b|o)$

$$P(b|o) = \frac{P(o|b)P(b)}{P(o)} = \frac{P(o|b)P(b)}{P(o|b)P(b) + P(o|r)P(r)} = 1/3$$

$P(B)$  prior probability - before we observe the fruit;  $P(B|F)$  - aposteriori probability - after we observe the fruit.

# Picking fruits. What is the probability that ...?

- ▶ red box: 2 apples, 6 oranges
- ▶ blue box: 3 apples, 1 orange



Scenario: Pick a box (say red box in 40% cases), then pick a fruit at random.

Given that we have chosen an orange, what is the probability that the box we chose was the blue one?

- A:  $1/4$
- B:  $3/5$
- C:  $1/3$
- D: Different value.

4 / 24

## Notes

Example serves for probability recap (sum, product rules, conditional probabilities, Bayes)

Random variables:

- Identity of the box  $B$ , two possible values  $r, b$
- Identity of the fruit  $F$ , two possible values  $a, o$

Info about picking a box  $P(B = r) = 0.4$  and  $P(B = b) = 0.6$ .

Conditional probabilities, given box selected:  $P(o|r) = 3/4$ ,  $P(a|r) = 1/4$ ,  $P(o|b) = 1/4$ ,  $P(a|b) = 3/4$ .

Answering questions:

- $P(F = a) = P(a|r)P(r) + P(a|b)P(b) = 11/20$
- $P(B = b|F = o) = P(b|o)$

$$P(b|o) = \frac{P(o|b)P(b)}{P(o)} = \frac{P(o|b)P(b)}{P(o|b)P(b) + P(o|r)P(r)} = 1/3$$

$P(B)$  prior probability - before we observe the fruit;  $P(B|F)$  - aposteriori probability - after we observe the fruit.

# Rules of probability and notation I

- ▶ **random variables**  $X, Y$
- ▶  $x_i$  where  $i = 1, \dots, M$  – values taken by variable  $X$
- ▶  $y_j$  where  $j = 1, \dots, L$  – values taken by variable  $Y$
- ▶  $P(X = x_i, Y = y_j)$  – probability that  $X$  takes the value  $x_i$  and  $Y$  takes  $y_j$  – **joint probability**
- ▶  $P(X = x_i)$  – probability that  $X$  takes the value  $x_i$
- ▶ **Sum rule of probability** :
  - ▶  $P(X = x_i) = \sum_{j=1}^L P(X = x_i, Y = y_j)$
  - ▶  $P(X = x_i)$  is sometimes called **marginal probability** – obtained by marginalizing / summing out the other variables
  - ▶ general rule, compact notation:  $P(X) = \sum_Y P(X, Y)$

---

## Notes

This and the following slides are just to formally recap what we learned when discussion boxes and fruits

# Rules of probability and notation II

- ▶ **Conditional probability** :  $P(Y = y_j | X = x_i)$
- ▶ **Product rule of probability** :
  - ▶  $P(X = x_i, Y = y_j) = P(Y = y_j | X = x_i)P(X = x_i)$
  - ▶ general rule, compact notation:  $P(X, Y) = P(Y|X)P(X)$
- ▶ **Bayes theorem** :
  - ▶ from  $P(X, Y) = P(Y, X)$  and product rule

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

$$\textit{posterior} = \frac{\textit{likelihood} \times \textit{prior}}{\textit{evidence}}$$

- ▶ **Independence** :  $P(X, Y) = P(X)P(Y)$

---

## Notes

What does it mean when we say that random variables  $X$  and  $Y$  are independent?



# Rules of probability and notation II

- ▶ **Conditional probability** :  $P(Y = y_j | X = x_i)$
- ▶ **Product rule of probability** :
  - ▶  $P(X = x_i, Y = y_j) = P(Y = y_j | X = x_i)P(X = x_i)$
  - ▶ general rule, compact notation:  $P(X, Y) = P(Y|X)P(X)$
- ▶ **Bayes theorem** :
  - ▶ from  $P(X, Y) = P(Y, X)$  and product rule

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

$$\textit{posterior} = \frac{\textit{likelihood} \times \textit{prior}}{\textit{evidence}}$$

- ▶ **Independence** :  $P(X, Y) = P(X)P(Y)$


---

## Notes

What does it mean when we say that random variables  $X$  and  $Y$  are independent?

# Boxes and Fruits: posterior? likelihood? prior? evidence?

$$\text{posterior} = \frac{\text{likelihood} \times \text{prior}}{\text{evidence}}$$



- ▶ posterior  
after observation
  - ▶ likelihood  
of an observation
  - ▶ prior  
before observation
  - ▶ evidence  
total observations
- ▶  $P(B)$
  - ▶  $P(F)$
  - ▶  $P(F | B)$
  - ▶  $P(B | F)$

## Notes

**B**oxes and **F**ruits: prior (before observation) -  $P(B)$ , likelihood (of observation) -  $P(F|B)$ , evidence (total observations)  $P(F)$ , posterior (after observation)  $P(B|F)$ .

Think about these terms, it helps to understand and remember.

## Decision example: Insure or not? (from late 1980s) [4]

A doctor calls: “Your HIV test is positive, 999/1000 you will die in 10 years. I’m sorry ...”.  
Insurance company does not want to insure a married couple.

- ▶ Was the doctor right?
- ▶ Was the insurance company rational?

What the doctor (and the company) knew:

- ▶ HIV test falsely positive only in 1 case out of 1000.
- ▶ Heterosexual male, has family, no drugs, no risk behavior.

---

### Notes

Equations/formulas are simple but not easy to (fully) understand.

Doctor:  $P(\text{positive test} \mid \text{healthy}) = \frac{1}{1000}$  but this is the *likelihood* which we learn before the patient's diagnosis (classification). More interesting and important is to know:  $P(\text{healthy} \mid \text{positive test})$  (*posterior*).

Think about 10000 samples of heterosexual males, family, ... . Statistically, there is just 1 HIV positive among them. Assume  $P(\text{negative test} \mid \text{infected}) \rightarrow 0$ . 1 person HIV positive will be tested positive, but also 10 other healthy persons will be tested positive. Hence  $P(\text{healthy} \mid \text{positive test}) = 10/11!$ . Or, for the doctor:  $P(\text{infected} \mid \text{positive test}) = \frac{1}{11}$  and not  $\frac{999}{1000}$ . The fact that a disease is rare matters a lot!

## Decision example: Insure or not? (from late 1980s) [4]

A doctor calls: “Your HIV test is positive, 999/1000 you will die in 10 years. I’m sorry ...”.  
Insurance company does not want to insure a married couple.

- ▶ Was the doctor right?
- ▶ Was the insurance company rational?

What the doctor (and the company) knew:

- ▶ HIV test falsely positive only in 1 case out of 1000.
- ▶ Heterosexual male, has family, no drugs, no risk behavior.

---

### Notes

Equations/formulas are simple but not easy to (fully) understand.

Doctor:  $P(\text{positive test} \mid \text{healthy}) = \frac{1}{1000}$  but this is the *likelihood* which we learn before the patient's diagnosis (classification). More interesting and important is to know:  $P(\text{healthy} \mid \text{positive test})$  (*posterior*).

Think about 10000 samples of heterosexual males, family, ... . Statistically, there is just 1 HIV positive among them. Assume  $P(\text{negative test} \mid \text{infected}) \rightarrow 0$ . 1 person HIV positive will be tested positive, but also 10 other healthy persons will be tested positive. Hence  $P(\text{healthy} \mid \text{positive test}) = 10/11!$ . Or, for the doctor:  $P(\text{infected} \mid \text{positive test}) = \frac{1}{11}$  and not  $\frac{999}{1000}$ . The fact that a disease is rare matters a lot!

## Decision example: Insure or not? (from late 1980s) [4]

A doctor calls: “Your HIV test is positive, 999/1000 you will die in 10 years. I’m sorry ...”. Insurance company does not want to insure a married couple.

- ▶ Was the doctor right?
- ▶ Was the insurance company rational?

What the doctor (and the company) knew:

- ▶ HIV test falsely positive only in 1 case out of 1000.
- ▶ Heterosexual male, has family, no drugs, no risk behavior.

---

### Notes

Equations/formulas are simple but not easy to (fully) understand.

Doctor:  $P(\text{positive test} \mid \text{healthy}) = \frac{1}{1000}$  but this is the *likelihood* which we learn before the patient's diagnosis (classification). More interesting and important is to know:  $P(\text{healthy} \mid \text{positive test})$  (*posterior*).

Think about 10000 samples of heterosexual males, family, .... Statistically, there is just 1 HIV positive among them. Assume  $P(\text{negative test} \mid \text{infected}) \rightarrow 0$ . 1 person HIV positive will be tested positive, but also 10 other healthy persons will be tested positive. Hence  $P(\text{healthy} \mid \text{positive test}) = 10/11!$ . Or, for the doctor:  $P(\text{infected} \mid \text{positive test}) = \frac{1}{11}$  and not  $\frac{999}{1000}$ . The fact that a disease is rare matters a lot!

## Decision example: Insure or not? (from late 1980s) [4]

A doctor calls: “Your HIV test is positive, 999/1000 you will die in 10 years. I’m sorry ...”. Insurance company does not want to insure a married couple.

- ▶ Was the doctor right?
- ▶ Was the insurance company rational?

What the doctor (and the company) knew:

- ▶ HIV test falsely positive only in 1 case out of 1000.
- ▶ Heterosexual male, has family, no drugs, no risk behavior.

---

### Notes

Equations/formulas are simple but not easy to (fully) understand.

Doctor:  $P(\text{positive test} \mid \text{healthy}) = \frac{1}{1000}$  but this is the *likelihood* which we learn before the patient's diagnosis (classification). More interesting and important is to know:  $P(\text{healthy} \mid \text{positive test})$  (*posterior*).

Think about 10000 samples of heterosexual males, family, ... Statistically, there is just 1 HIV positive among them. Assume  $P(\text{negative test} \mid \text{infected}) \rightarrow 0$ . 1 person HIV positive will be tested positive, but also 10 other healthy persons will be tested positive. Hence  $P(\text{healthy} \mid \text{positive test}) = 10/11!$ . Or, for the doctor:  $P(\text{infected} \mid \text{positive test}) = \frac{1}{11}$  and not  $\frac{999}{1000}$ . The fact that a disease is rare matters a lot!

## Decision example: Insure or not? (from late 1980s) [4]

A doctor calls: “Your HIV test is positive, 999/1000 you will die in 10 years. I’m sorry ...”. Insurance company does not want to insure a married couple.

- ▶ Was the doctor right?
- ▶ Was the insurance company rational?

What the doctor (and the company) knew:

- ▶ HIV test falsely positive only in 1 case out of 1000.
- ▶ Heterosexual male, has family, no drugs, no risk behavior.

---

### Notes

Equations/formulas are simple but not easy to (fully) understand.

Doctor:  $P(\text{positive test} \mid \text{healthy}) = \frac{1}{1000}$  but this is the *likelihood* which we learn before the patient's diagnosis (classification). More interesting and important is to know:  $P(\text{healthy} \mid \text{positive test})$  (*posterior*).

Think about 10000 samples of heterosexual males, family, ... Statistically, there is just 1 HIV positive among them. Assume  $P(\text{negative test} \mid \text{infected}) \rightarrow 0$ . 1 person HIV positive will be tested positive, but also 10 other healthy persons will be tested positive. Hence  $P(\text{healthy} \mid \text{positive test}) = 10/11!$ . Or, for the doctor:  $P(\text{infected} \mid \text{positive test}) = \frac{1}{11}$  and not  $\frac{999}{1000}$ . The fact that a disease is rare matters a lot!

## Decision: guilty or not? (people of CA vs Collins, 1968) [4]

- ▶ Robbery, LA 1964, fuzzy evidence of the offenders:
  - ▶ female, around 65 kg
  - ▶ wearing something dark
  - ▶ hairs of light color, between light and dark blond, in a ponytail
- ▶ At the same time, additional evidence close to the crime scene:
  - ▶ loud scream, yelling, looking at the this direction
  - ...
  - ▶ a woman sitting into a yellow car
  - ▶ car starts immediately and passes close to the additional witness
  - ▶ a black man with beard and moustache was driving
- ▶ No more evidence
- ▶ Testimony of both the victim and the witness not unambiguous (didn't recognize suspects)
- ▶ Still, the suspects were sentenced to jail.

9 / 24

---

### Notes

Wrong use of independence assumption:

$$\begin{aligned}P(\text{yellow car}) &= 1/10 \\P(\text{man with moustache}) &= 1/4 \\P(\text{black man with beard}) &= 1/10 \\P(\text{woman with pony tail}) &= 1/10 \\P(\text{woman blond hair}) &= 1/3 \\P(\text{mix race pair in a car}) &= 1/1000\end{aligned}$$

and mistakenly confusing probability


$$P(\text{randomly selected pair matches discussed characteristics})$$

giving  $P = 1/12000000$   
with the needed:

$$P(\text{a pair matching characteristics is guilty})$$



# Classification example: What's the fish?



- ▶ Factory for fish processing
- ▶ 2 classes  $s_{1,2}$ :
  - ▶ salmon
  - ▶ sea bass
- ▶ Features  $\vec{x}$ : length, width, lightness etc. from a camera

## Notes

- Sea (European) bass, [https://en.wikipedia.org/wiki/European\\_bass](https://en.wikipedia.org/wiki/European_bass). In czech it is Mořčák evropský or Mořský vlk.
- Salmon, <https://en.wikipedia.org/wiki/Salmon>, is Losos in Czech.

# Fish – classification using probability

$$posterior = \frac{likelihood \times prior}{evidence}$$

## ▶ Notation for classification problem

- ▶ Classes  $s_j \in S$  (e.g., salmon, sea bass)
- ▶ Features  $x_i \in X$  or feature vectors  $(\vec{x}_i)$  (also called attributes)

## ▶ Optimal classification of $\vec{x}$ :

$$\delta^*(\vec{x}) = \arg \max_j P(s_j | \vec{x})$$

- ▶ We thus choose the most probable class for a given feature vector.
- ▶ Both likelihood and prior are taken into account – recall Bayes rule:

$$P(s_j | \vec{x}) = \frac{P(\vec{x} | s_j) P(s_j)}{P(\vec{x})}$$

## ▶ Can we do (classify) better?

11 / 24

---

### Notes

Assuming we know the true  $P(\vec{x} | s_j)$ ,  $P(s_j)$ ,  $P(\vec{x})$  we *cannot* do better! Bayesian decision is the optimal one!

# Fish – classification using probability

$$posterior = \frac{likelihood \times prior}{evidence}$$

- ▶ Notation for classification problem
  - ▶ Classes  $s_j \in S$  (e.g., salmon, sea bass)
  - ▶ Features  $x_i \in X$  or feature vectors  $(\vec{x}_i)$  (also called attributes)
- ▶ Optimal classification of  $\vec{x}$ :

$$\delta^*(\vec{x}) = \arg \max_j P(s_j | \vec{x})$$

- ▶ We thus choose the **most probable class for a given feature vector** .
- ▶ Both likelihood and prior are taken into account – recall Bayes rule:

$$P(s_j | \vec{x}) = \frac{P(\vec{x} | s_j) P(s_j)}{P(\vec{x})}$$

- ▶ Can we do (classify) better?

11 / 24

---

## Notes

Assuming we know the true  $P(\vec{x} | s_j)$ ,  $P(s_j)$ ,  $P(\vec{x})$  we *cannot* do better! Bayesian decision is the optimal one!

# Bayes classification in practice

- ▶ Usually we are not given  $P(s|\vec{x})$ 
  - ▶ It has to be estimated from already classified examples – training data
  - ▶ For discrete  $\vec{x}$ , training examples  $(\vec{x}_1, s_1), (\vec{x}_2, s_2), \dots, (\vec{x}_l, s_l)$ 
    - ▶ so-called i.i.d (independent, identically distributed) multiset
    - ▶ every  $(\vec{x}, s)$  is drawn independently from  $P(\vec{x}, s)$
  - ▶ Without knowing anything about the distribution, a non-parametric estimate:

$$P(s|\vec{x}) \approx \frac{\# \text{ examples where } \vec{x}_i = \vec{x} \text{ and } s_j = s}{\# \text{ examples where } \vec{x}_i = \vec{x}}$$

- ▶ Hard in practice:
  - ▶ To reliably estimate  $P(s|\vec{x})$ , the number of examples grows exponentially with the number of elements of  $\vec{x}$ .
    - ▶ e.g. with the number of pixels in images
    - ▶ curse of dimensionality
    - ▶ denominator often 0

12 / 24

---

## Notes

Why hard? Way too many various  $\vec{x}$ . Think about simple binary  $10 \times 10$  image -  $\vec{x}$  contains 0, 1, position matters. What is the total number of unique images? Think binary,  $1 \times 8$  binary image? It is very hard almost impossible to sample – collect training data – joint probability distribution.

# Bayes classification in practice

- ▶ Usually we are not given  $P(s|\vec{x})$
- ▶ It has to be estimated from already classified examples – training data
- ▶ For discrete  $\vec{x}$ , training examples  $(\vec{x}_1, s_1), (\vec{x}_2, s_2), \dots, (\vec{x}_l, s_l)$ 
  - ▶ so-called i.i.d (independent, identically distributed) multiset
  - ▶ every  $(\vec{x}_i, s_i)$  is drawn independently from  $P(\vec{x}, s)$
- ▶ Without knowing anything about the distribution, a non-parametric estimate:

$$P(s|\vec{x}) \approx \frac{\# \text{ examples where } \vec{x}_i = \vec{x} \text{ and } s_i = s}{\# \text{ examples where } \vec{x}_i = \vec{x}}$$

- ▶ Hard in practice:
  - ▶ To reliably estimate  $P(s|\vec{x})$ , the number of examples grows exponentially with the number of elements of  $\vec{x}$ .
    - ▶ e.g. with the number of pixels in images
    - ▶ curse of dimensionality
    - ▶ denominator often 0

12 / 24

---

## Notes

Why hard? Way too many various  $\vec{x}$ . Think about simple binary  $10 \times 10$  image -  $\vec{x}$  contains 0, 1, position matters. What is the total number of unique images? Think binary,  $1 \times 8$  binary image? It is very hard almost impossible to sample – collect training data – joint probability distribution.

# Bayes classification in practice

- ▶ Usually we are not given  $P(s|\vec{x})$
- ▶ It has to be estimated from already classified examples – training data
- ▶ For discrete  $\vec{x}$ , training examples  $(\vec{x}_1, s_1), (\vec{x}_2, s_2), \dots, (\vec{x}_l, s_l)$ 
  - ▶ so-called i.i.d (independent, identically distributed) multiset
  - ▶ every  $(\vec{x}_i, s_i)$  is drawn independently from  $P(\vec{x}, s)$
- ▶ Without knowing anything about the distribution, a non-parametric estimate:

$$P(s|\vec{x}) \approx \frac{\# \text{ examples where } \vec{x}_i = \vec{x} \text{ and } s_i = s}{\# \text{ examples where } \vec{x}_i = \vec{x}}$$

- ▶ Hard in practice:
  - ▶ To reliably estimate  $P(s|\vec{x})$ , the number of examples grows exponentially with the number of elements of  $\vec{x}$ .
    - ▶ e.g. with the number of pixels in images
    - ▶ curse of dimensionality
    - ▶ denominator often 0

---

## Notes

Why hard? Way too many various  $\vec{x}$ . Think about simple binary  $10 \times 10$  image -  $\vec{x}$  contains 0, 1, position matters. What is the total number of unique images? Think binary,  $1 \times 8$  binary image? It is very hard almost impossible to sample – collect training data – joint probability distribution.

# Naïve Bayes classification

- ▶ For efficient classification we must thus rely on additional assumptions.
- ▶ In the exceptional case of **statistical independence** between  $\vec{x}$  components for each class  $s$  it holds

$$P(\vec{x}|s) = P(x[1]|s) \cdot P(x[2]|s) \cdot \dots$$

- ▶ Use simple Bayes law and maximize:

$$P(s|\vec{x}) = \frac{P(\vec{x}|s)P(s)}{P(\vec{x})} = \frac{P(s)}{P(\vec{x})} P(x[1]|s) \cdot P(x[2]|s) \cdot \dots =$$

- ▶ No combinatorial curse in estimating  $P(s)$  and  $P(x[i]|s)$  separately for each  $i$  and  $s$ .
- ▶ No need to estimate  $P(\vec{x})$ . (Why?)
- ▶  $P(s)$  may be provided apriori.
- ▶ **naïve** = when used despite statistical dependence

13 / 24

---

## Notes

Why naïve at all? Consider  $N$ -dimensional space,  $8$ -bit values. Instead of problem  $8^N$  we have  $8 \times N$  problem. Think about statistical independence. Example1: person's weight and height. Are they independent? Example2: pixel values in images.

We will talk about learning classifiers more in next lectures.

# Decision making under uncertainty

- ▶ An important feature of intelligent systems
  - ▶ make the best possible decision
  - ▶ in uncertain conditions.
- ▶ Example: Take a tram OR subway from *A* to *B*?
  - ▶ Tram: timetables imply a quicker route, but adherence uncertain.
  - ▶ Subway: longer route, but adherence almost certain.
- ▶ Example: where to route a letter with this ZIP?

▶ 15700? 15706? 15200? 15206?

- ▶ What is the optimal decision ?
- ▶ Both examples fall into the same framework.

14 / 24

---

## Notes

There are *costs* associated with a decision. E.g. at fish packing plant, customers may not mind so much if some pieces of salmon end up in sea bass cans, but they will be protesting if the opposite happens. So making an error “one way” has higher cost than “the other way”. This impacts where decision boundaries for classification should optimally be drawn.



# Decision making under uncertainty

- ▶ An important feature of intelligent systems
  - ▶ make the best possible decision
  - ▶ in uncertain conditions.
- ▶ **Example:** Take a tram OR subway from *A* to *B*?
  - ▶ Tram: timetables imply a quicker route, but adherence uncertain.
  - ▶ Subway: longer route, but adherence almost certain.
- ▶ Example: where to route a letter with this ZIP?

▶ 15700? 15706? 15200? 15206?

- ▶ What is the optimal decision ?
- ▶ Both examples fall into the same framework.

14 / 24

---

## Notes

There are *costs* associated with a decision. E.g. at fish packing plant, customers may not mind so much if some pieces of salmon end up in sea bass cans, but they will be protesting if the opposite happens. So making an error “one way” has higher cost than “the other way”. This impacts where decision boundaries for classification should optimally be drawn.

# Decision making under uncertainty

- ▶ An important feature of intelligent systems
  - ▶ make the best possible decision
  - ▶ in uncertain conditions.
- ▶ **Example:** Take a tram OR subway from *A* to *B*?
  - ▶ Tram: timetables imply a quicker route, but adherence uncertain.
  - ▶ Subway: longer route, but adherence almost certain.
- ▶ **Example:** where to route a letter with this ZIP?

A handwritten ZIP code '15700' is shown. The '1' is on the left, followed by '57' and '00'. The '5' and '7' are connected, and the '00' is written as two loops.

- ▶ 15700? 15706? 15200? 15206?
- ▶ What is the optimal decision?
- ▶ Both examples fall into the same framework.


---

## Notes

There are *costs* associated with a decision. E.g. at fish packing plant, customers may not mind so much if some pieces of salmon end up in sea bass cans, but they will be protesting if the opposite happens. So making an error “one way” has higher cost than “the other way”. This impacts where decision boundaries for classification should optimally be drawn.

# Decision making under uncertainty

- ▶ An important feature of intelligent systems
  - ▶ make the best possible decision
  - ▶ in uncertain conditions.
- ▶ **Example:** Take a tram OR subway from *A* to *B*?
  - ▶ Tram: timetables imply a quicker route, but adherence uncertain.
  - ▶ Subway: longer route, but adherence almost certain.
- ▶ **Example:** where to route a letter with this ZIP?

A handwritten ZIP code '15700' is shown. The '1' is on the left, followed by '57' and '00'. The '00' is written with two loops. There is a faint dotted line between '57' and '00'.

- ▶ 15700? 15706? 15200? 15206?
- ▶ What is the **optimal decision** ?
- ▶ Both examples fall into the same framework.

---

## Notes

There are *costs* associated with a decision. E.g. at fish packing plant, customers may not mind so much if some pieces of salmon end up in sea bass cans, but they will be protesting if the opposite happens. So making an error “one way” has higher cost than “the other way”. This impacts where decision boundaries for classification should optimally be drawn.

## Example: What to cook for a dinner [3]

- ▶ Wife coming back from work. Husband: what to cook for dinner?
- ▶ 3 dishes ( decisions ) in his repertoire:
  - ▶ *nothing* ... don't bother cooking  $\Rightarrow$  no work but makes wife upset
  - ▶ *pizza* ... microwave a frozen pizza  $\Rightarrow$  not much work but won't impress
  - ▶ *g.T.c.* ... general Tso's chicken  $\Rightarrow$  will make her day, but very laborious.
- ▶ Hassle incurred by the individual options depends wife's feeling
- ▶ For each of the 9 possible situation (3 possible decisions  $\times$  3 possible states) the hassle is quantified by a loss function  $l(d, s)$ :

| $l(s, d)$              | $d = \textit{nothing}$ | $d = \textit{pizza}$ | $d = \textit{g.T.c.}$ |
|------------------------|------------------------|----------------------|-----------------------|
| $s = \textit{good}$    | 0                      | 2                    | 4                     |
| $s = \textit{average}$ | 5                      | 3                    | 5                     |
| $s = \textit{bad}$     | 10                     | 9                    | 6                     |

Wife's state of mind is an uncertain state.

---

### Notes

Was the state known the decision would be simple.

## Example: What to cook for a dinner [3]

- ▶ Wife coming back from work. Husband: what to cook for dinner?
- ▶ 3 dishes ( **decisions** ) in his repertoire:
  - ▶ *nothing* ... **don't bother cooking**  $\Rightarrow$  no work but makes wife upset
  - ▶ *pizza* ... **microwave a frozen pizza**  $\Rightarrow$  not much work but won't impress
  - ▶ *g.T.c.* ... **general Tso's chicken**  $\Rightarrow$  will make her day, but very laborious.
- ▶ Hassle incurred by the individual options depends wife's feeling
- ▶ For each of the 9 possible situation (3 possible decisions  $\times$  3 possible states) the hassle is quantified by a loss function  $l(d, s)$ :

| $l(s, d)$              | $d = \textit{nothing}$ | $d = \textit{pizza}$ | $d = \textit{g.T.c.}$ |
|------------------------|------------------------|----------------------|-----------------------|
| $s = \textit{good}$    | 0                      | 2                    | 4                     |
| $s = \textit{average}$ | 5                      | 3                    | 5                     |
| $s = \textit{bad}$     | 10                     | 9                    | 6                     |

Wife's state of mind is an uncertain state.

---

### Notes

Was the state known the decision would be simple.

## Example: What to cook for a dinner [3]

- ▶ Wife coming back from work. Husband: what to cook for dinner?
- ▶ 3 dishes ( **decisions** ) in his repertoire:
  - ▶ *nothing* ... **don't bother cooking**  $\Rightarrow$  no work but makes wife upset
  - ▶ *pizza* ... **microwave a frozen pizza**  $\Rightarrow$  not much work but won't impress
  - ▶ *g.T.c.* ... **general Tso's chicken**  $\Rightarrow$  will make her day, but very laborious.
- ▶ Hassle incurred by the individual options depends wife's feeling
- ▶ For each of the 9 possible situation (3 possible decisions  $\times$  3 possible states) the hassle is quantified by a **loss function**  $l(d, s)$ :

| $l(s, d)$              | $d = \textit{nothing}$ | $d = \textit{pizza}$ | $d = \textit{g.T.c.}$ |
|------------------------|------------------------|----------------------|-----------------------|
| $s = \textit{good}$    | 0                      | 2                    | 4                     |
| $s = \textit{average}$ | 5                      | 3                    | 5                     |
| $s = \textit{bad}$     | 10                     | 9                    | 6                     |

Wife's state of mind is an uncertain state.

---

### Notes

Was the state known the decision would be simple.

## Example: What to cook for a dinner [3]

- ▶ Wife coming back from work. Husband: what to cook for dinner?
- ▶ 3 dishes ( **decisions** ) in his repertoire:
  - ▶ *nothing* ... **don't bother cooking**  $\Rightarrow$  no work but makes wife upset
  - ▶ *pizza* ... **microwave a frozen pizza**  $\Rightarrow$  not much work but won't impress
  - ▶ *g.T.c.* ... **general Tso's chicken**  $\Rightarrow$  will make her day, but very laborious.
- ▶ Hassle incurred by the individual options depends wife's feeling
- ▶ For each of the 9 possible situation (3 possible decisions  $\times$  3 possible states) the hassle is quantified by a **loss function**  $l(d, s)$ :

| $l(s, d)$              | $d = \textit{nothing}$ | $d = \textit{pizza}$ | $d = \textit{g.T.c.}$ |
|------------------------|------------------------|----------------------|-----------------------|
| $s = \textit{good}$    | 0                      | 2                    | 4                     |
| $s = \textit{average}$ | 5                      | 3                    | 5                     |
| $s = \textit{bad}$     | 10                     | 9                    | 6                     |

Wife's state of mind is an **uncertain state**.

---

### Notes

Was the state known the decision would be simple.

## Example (cont'd), State uncertain, ...

- ▶ Husband's experiment: He tells her he accidentally overtaped their wedding video and observes her reaction
- ▶ Anticipates 4 possible reactions:
  - ▶ *mild* ... all right, we keep our memories.
  - ▶ *irritated* ... how many times do I have to tell you...
  - ▶ *upset* ... Why did I marry this guy?
  - ▶ *alarming* ... silence
- ▶ The reaction is a measurable attribute ( "feature" ) of the mind state.
- ▶ From experience, the husband knows how individual reactions are probable in each state of mind; this is captured by the joint distribution  $P(x, s)$  .

| $P(x, s)$     | $x = mild$ | $x = irritated$ | $x = upset$ | $x = alarming$ |
|---------------|------------|-----------------|-------------|----------------|
| $s = good$    | 0.35       | 0.28            | 0.07        | 0.00           |
| $s = average$ | 0.04       | 0.10            | 0.04        | 0.02           |
| $s = bad$     | 0.00       | 0.02            | 0.05        | 0.03           |

16 / 24

---

### Notes

Joint distribution. Husband tried similar experiment multiple times, gathered some evidence ...

Instead of complicated experiment with overtaping wedding video, think about asking "when are you coming home?" .



## Example (cont'd), State uncertain, ...

- ▶ Husband's experiment. He tells her he accidentally overtaped their wedding video and observes her reaction
- ▶ Anticipates 4 possible reactions:
  - ▶ *mild* ... all right, we keep our memories.
  - ▶ *irritated* ... how many times do I have to tell you...
  - ▶ *upset* ... Why did I marry this guy?
  - ▶ *alarming* ... silence
- ▶ The reaction is a measurable attribute ( "feature" ) of the mind state.
- ▶ From experience, the husband knows how individual reactions are probable in each state of mind; this is captured by the joint distribution  $P(x, s)$  .

| $P(x, s)$     | $x = mild$ | $x = irritated$ | $x = upset$ | $x = alarming$ |
|---------------|------------|-----------------|-------------|----------------|
| $s = good$    | 0.35       | 0.28            | 0.07        | 0.00           |
| $s = average$ | 0.04       | 0.10            | 0.04        | 0.02           |
| $s = bad$     | 0.00       | 0.02            | 0.05        | 0.03           |

16 / 24

---

### Notes

Joint distribution. Husband tried similar experiment multiple times, gathered some evidence ...

Instead of complicated experiment with overtaping wedding video, think about asking "when are you coming home?" .

## Example (cont'd), State uncertain, ...

- ▶ Husband's experiment. He tells her he accidentally overtaped their wedding video and observes her reaction
- ▶ Anticipates 4 possible reactions:
  - ▶ *mild* ... all right, we keep our memories.
  - ▶ *irritated* ... how many times do I have to tell you....
  - ▶ *upset* ... Why did I marry this guy?
  - ▶ *alarming* ... silence
- ▶ The reaction is a measurable **attribute** ( **"feature"** ) of the mind state.
- ▶ From experience, the husband knows how individual reactions are probable in each state of mind; this is captured by the joint distribution  $P(x, s)$ .

| $P(x, s)$     | $x = mild$ | $x = irritated$ | $x = upset$ | $x = alarming$ |
|---------------|------------|-----------------|-------------|----------------|
| $s = good$    | 0.35       | 0.28            | 0.07        | 0.00           |
| $s = average$ | 0.04       | 0.10            | 0.04        | 0.02           |
| $s = bad$     | 0.00       | 0.02            | 0.05        | 0.03           |

16 / 24

---

### Notes

Joint distribution. Husband tried similar experiment multiple times, gathered some evidence ...

Instead of complicated experiment with overtaping wedding video, think about asking "when are you coming home?".

## Example (cont'd), State uncertain, ...

- ▶ Husband's experiment. He tells her he accidentally overtaped their wedding video and observes her reaction
- ▶ Anticipates 4 possible reactions:
  - ▶ *mild* ... all right, we keep our memories.
  - ▶ *irritated* ... how many times do I have to tell you....
  - ▶ *upset* ... Why did I marry this guy?
  - ▶ *alarming* ... silence
- ▶ The reaction is a measurable **attribute** ( "**feature**" ) of the mind state.
- ▶ From experience, the husband knows how individual reactions are probable in each state of mind; this is captured by the **joint distribution**  $P(x, s)$  .

| $P(x, s)$     | $x = mild$ | $x = irritated$ | $x = upset$ | $x = alarming$ |
|---------------|------------|-----------------|-------------|----------------|
| $s = good$    | 0.35       | 0.28            | 0.07        | 0.00           |
| $s = average$ | 0.04       | 0.10            | 0.04        | 0.02           |
| $s = bad$     | 0.00       | 0.02            | 0.05        | 0.03           |

---

### Notes

Joint distribution. Husband tried similar experiment multiple times, gathered some evidence ...

Instead of complicated experiment with overtaping wedding video, think about asking "when are you coming home?" .

# Decision strategy

- ▶ **Decision strategy** : a rule selecting a decision for *any given value* of the measured attribute(s).
- ▶ i.e. function  $d = \delta(x)$ .
- ▶ Example of husband's possible strategies:

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_4(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>nothing</i>     | <i>nothing</i>        |

- ▶ How many strategies?
- ▶ How to define which strategy is best? How to sort them by quality?
- ▶ Define the risk of a strategy as a mean (expected) loss value .

$$r(\delta) = \sum_x \sum_s l(s, \delta(x)) P(x, s)$$

---

## Notes

Overall,  $3^4 = 81$  possible strategies (3 possible decisions for each of the 4 possible attribute values). There is some analogy of states and possible actions. Here, we reason about states - which are 3 (state of mind) - from features which are 4.

Any given value (of measured attribute) ... Think about any possible state

# Decision strategy

- ▶ **Decision strategy** : a rule selecting a decision for *any given value* of the measured attribute(s).
- ▶ i.e. function  $d = \delta(x)$ .
- ▶ Example of husband's possible strategies:

| $\delta(x)$     | $x = mild$     | $x = irritated$ | $x = upset$    | $x = alarming$ |
|-----------------|----------------|-----------------|----------------|----------------|
| $\delta_1(x) =$ | <i>nothing</i> | <i>nothing</i>  | <i>pizza</i>   | <i>g.T.c.</i>  |
| $\delta_2(x) =$ | <i>nothing</i> | <i>pizza</i>    | <i>g.T.c.</i>  | <i>g.T.c.</i>  |
| $\delta_3(x) =$ | <i>g.T.c.</i>  | <i>g.T.c.</i>   | <i>g.T.c.</i>  | <i>g.T.c.</i>  |
| $\delta_4(x) =$ | <i>nothing</i> | <i>nothing</i>  | <i>nothing</i> | <i>nothing</i> |

- ▶ How many strategies?
  - ▶ How to define which strategy is best? How to sort them by quality?
  - ▶ Define the *risk* of a strategy as a mean (expected) loss value . . .

$$r(\delta) = \sum_x \sum_s l(s, \delta(x)) P(s; x)$$

---

## Notes

Overall,  $3^4 = 81$  possible strategies (3 possible decisions for each of the 4 possible attribute values). There is some analogy of states and possible actions. Here, we reason about states - which are 3 (state of mind) - from features which are 4.

Any given value (of measured attribute) . . . Think about any possible state

# Decision strategy

- ▶ **Decision strategy** : a rule selecting a decision for *any given value* of the measured attribute(s).
- ▶ i.e. function  $d = \delta(x)$ .
- ▶ Example of husband's possible strategies:

| $\delta(x)$     | $x = mild$     | $x = irritated$ | $x = upset$    | $x = alarming$ |
|-----------------|----------------|-----------------|----------------|----------------|
| $\delta_1(x) =$ | <i>nothing</i> | <i>nothing</i>  | <i>pizza</i>   | <i>g.T.c.</i>  |
| $\delta_2(x) =$ | <i>nothing</i> | <i>pizza</i>    | <i>g.T.c.</i>  | <i>g.T.c.</i>  |
| $\delta_3(x) =$ | <i>g.T.c.</i>  | <i>g.T.c.</i>   | <i>g.T.c.</i>  | <i>g.T.c.</i>  |
| $\delta_4(x) =$ | <i>nothing</i> | <i>nothing</i>  | <i>nothing</i> | <i>nothing</i> |

- ▶ How many strategies?
- ▶ How to define which strategy is best? How to sort them by quality?
- ▶ Define the **risk of a strategy** as a **mean (expected) loss value** .

$$r(\delta) = \sum_x \sum_s l(s, \delta(x)) P(x, s)$$

---

## Notes

Overall,  $3^4 = 81$  possible strategies (3 possible decisions for each of the 4 possible attribute values). There is some analogy of states and possible actions. Here, we reason about states - which are 3 (state of mind) - from features which are 4.

Any given value (of measured attribute) ... Think about any possible state

$$\text{Calculating } r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$$

| $l(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |
|----------------------|----------------------|--------------------|---------------------|
| $s = \text{good}$    | 0                    | 2                  | 4                   |
| $s = \text{average}$ | 5                    | 3                  | 5                   |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\vdots$        | $\vdots$          | $\vdots$               | $\vdots$           | $\vdots$              |

Do we need to evaluate all possible strategies?  $P(x, s) = P(s|x)P(x)$

### Notes

Risk depends on strategy (decisions). Strategy (decisions) depends on observation. Loss comes from decision and state. The total weighted average is weighted by joint probability of observation and state.

Calculate  $r(\delta_1)$  and  $r(\delta_2)$ , which strategy is better?

$$\text{Calculating } r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$$

| $l(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |
|----------------------|----------------------|--------------------|---------------------|
| $s = \text{good}$    | 0                    | 2                  | 4                   |
| $s = \text{average}$ | 5                    | 3                  | 5                   |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\vdots$        | $\vdots$          | $\vdots$               | $\vdots$           | $\vdots$              |

Do we need to evaluate all possible strategies?  $P(x, s) = P(s|x)P(x)$

### Notes

Risk depends on strategy (decisions). Strategy (decisions) depends on observation. Loss comes from decision and state. The total weighted average is weighted by joint probability of observation and state.

Calculate  $r(\delta_1)$  and  $r(\delta_2)$ , which strategy is better?



# Calculating $r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$

| $l(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |
|----------------------|----------------------|--------------------|---------------------|
| $s = \text{good}$    | 0                    | 2                  | 4                   |
| $s = \text{average}$ | 5                    | 3                  | 5                   |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\vdots$        | $\vdots$          | $\vdots$               | $\vdots$           | $\vdots$              |

Do we need to evaluate all possible strategies?  $P(x, s) = P(s|x)P(x)$

## Notes

Risk depends on strategy (decisions). Strategy (decisions) depends on observation. Loss comes from decision and state. The total weighted average is weighted by joint probability of observation and state.

Calculate  $r(\delta_1)$  and  $r(\delta_2)$ , which strategy is better?

$$\text{Calculating } r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$$

| $l(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |
|----------------------|----------------------|--------------------|---------------------|
| $s = \text{good}$    | 0                    | 2                  | 4                   |
| $s = \text{average}$ | 5                    | 3                  | 5                   |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\vdots$        | $\vdots$          | $\vdots$               | $\vdots$           | $\vdots$              |

Do we need to evaluate all possible strategies?

$$P(x, s) = P(s|x)P(x)$$

### Notes

Risk depends on strategy (decisions). Strategy (decisions) depends on observation. Loss comes from decision and state. The total weighted average is weighted by joint probability of observation and state.

Calculate  $r(\delta_1)$  and  $r(\delta_2)$ , which strategy is better?

$$\text{Calculating } r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$$

| $l(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |  |
|----------------------|----------------------|--------------------|---------------------|--|
| $s = \text{good}$    | 0                    | 2                  | 4                   |  |
| $s = \text{average}$ | 5                    | 3                  | 5                   |  |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |  |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta_1(x) =$ | <i>nothing</i>    | <i>nothing</i>         | <i>pizza</i>       | <i>g.T.c.</i>         |
| $\delta_2(x) =$ | <i>nothing</i>    | <i>pizza</i>           | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\delta_3(x) =$ | <i>g.T.c.</i>     | <i>g.T.c.</i>          | <i>g.T.c.</i>      | <i>g.T.c.</i>         |
| $\vdots$        | $\vdots$          | $\vdots$               | $\vdots$           | $\vdots$              |

Do we need to evaluate all possible strategies?  $P(x, s) = P(s|x)P(x)$

### Notes

Risk depends on strategy (decisions). Strategy (decisions) depends on observation. Loss comes from decision and state. The total weighted average is weighted by joint probability of observation and state.

Calculate  $r(\delta_1)$  and  $r(\delta_2)$ , which strategy is better?

## Bayes optimal strategy

- ▶ The **Bayes optimal strategy** : one minimizing mean risk.

$$\delta^* = \arg \min_{\delta} r(\delta)$$

- ▶ From  $P(x, s) = P(s|x)P(x)$  (Bayes rule), we have

$$\begin{aligned} r(\delta) &= \sum_x \sum_s l(s, \delta(x)) P(x, s) = \sum_s \sum_x l(s, \delta(x)) P(s|x) P(x) \\ &= \sum_x P(x) \underbrace{\sum_s l(s, \delta(x)) P(s|x)}_{\text{Conditional risk}} \end{aligned}$$

- ▶ The optimal strategy is obtained by minimizing the conditional risk *separately* for each  $x$ :

$$\delta^*(x) = \arg \min_d \sum_s l(s, d) P(s|x)$$

# Optimal strategy: $\delta^*(x) = \arg \min_d \sum_s I(s, d)P(s|x)$

| $I(s, d)$            | $d = \text{nothing}$ | $d = \text{pizza}$ | $d = \text{g.T.c.}$ |
|----------------------|----------------------|--------------------|---------------------|
| $s = \text{good}$    | 0                    | 2                  | 4                   |
| $s = \text{average}$ | 5                    | 3                  | 5                   |
| $s = \text{bad}$     | 10                   | 9                  | 6                   |

| $P(x, s)$            | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.35              | 0.28                   | 0.07               | 0.00                  |
| $s = \text{average}$ | 0.04              | 0.10                   | 0.04               | 0.02                  |
| $s = \text{bad}$     | 0.00              | 0.02                   | 0.05               | 0.03                  |

| $\delta(x)$     | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|-----------------|-------------------|------------------------|--------------------|-----------------------|
| $\delta^*(x) =$ | ??                | ??                     | ??                 | ??                    |

## Notes

We need to recompute the table of joint probability  $P(s, x)$  into table of conditional probabilities  $P(s|x)$ .

This can be done in two ways. A: Using product rule,  $P(s|x) = P(s, x)/P(x)$ .

First, to get  $P(x)$ , we use Sum rule (marginalizing).

| $P(x)$ | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|--------|-------------------|------------------------|--------------------|-----------------------|
|        | 0.39              | 0.40                   | 0.16               | 0.05                  |

Second, applying product rule,  $P(s|x) = P(s, x)/P(x)$ .

B: calculating the probability on a "per column basis".

E.g. for the first cell, A:  $0.35/0.39 = 0.897$  B:  $0.35/(0.35 + 0.04)$

| $P(s x)$             | $x = \text{mild}$ | $x = \text{irritated}$ | $x = \text{upset}$ | $x = \text{alarming}$ |
|----------------------|-------------------|------------------------|--------------------|-----------------------|
| $s = \text{good}$    | 0.897             | 0.7                    | 0.438              | 0.00                  |
| $s = \text{average}$ | 0.103             | 0.25                   | 0.25               | 0.4                   |
| $s = \text{bad}$     | 0.00              | 0.125                  | 0.313              | 0.6                   |

Having the table of all  $P(s|x)$  we just

mechanically insert into the equation in the slide title.

# Statistical decision making: wrapping up

## ▶ Given:

- ▶ A set of possible **states** :  $\mathcal{S}$
- ▶ A set of possible **decisions** :  $\mathcal{D}$
- ▶ A **loss function**  $l : \mathcal{D} \times \mathcal{S} \rightarrow \mathfrak{R}$
- ▶ The range  $\mathcal{X}$  of the **attribute**
- ▶ Distribution  $P(x, s)$ ,  $x \in \mathcal{X}$ ,  $s \in \mathcal{S}$ .

## ▶ Define:

- ▶ **Strategy** : function  $\delta : \mathcal{X} \rightarrow \mathcal{D}$
- ▶ **Risk of strategy**  $\delta : r(\delta) = \sum_x \sum_s l(s, \delta(x))P(x, s)$

## ▶ Bayes problem:

- ▶ Goal: find the optimal strategy  $\delta^* = \arg \min_{\delta \in \Delta} r(\delta)$
- ▶ Solution:  $\delta^*(x) = \arg \min_d \sum_s l(s, d)P(s|x)$

## A special case - Bayesian *classification*

- ▶ Bayesian classification is a special case of statistical decision theory:
  - ▶ Attribute vector  $\vec{x} = (x_1, x_2, \dots)$ : pixels 1, 2, ...
  - ▶ **State set  $\mathcal{S}$  = decision set  $\mathcal{D} = \{0, 1, \dots, 9\}$ .**
  - ▶ **State = actual class, Decision = recognized class**
  - ▶ Loss function:

$$l(s, d) = \begin{cases} 0, & d = s \\ 1, & d \neq s \end{cases}$$

$$\delta^*(\vec{x}) = \arg \min_d \sum_s \underbrace{l(s, d)}_{0 \text{ if } d=s} P(s|\vec{x}) = \arg \min_d \sum_{s \neq d} P(s|\vec{x})$$

Obviously  $\sum_s P(s|\vec{x}) = 1$ , then:

$$P(d|\vec{x}) + \sum_{s \neq d} P(s|\vec{x}) = 1$$

Inserting into above:

$$\delta^*(\vec{x}) = \arg \min_d [1 - P(d|\vec{x})] = \arg \max_d P(d|\vec{x})$$

---

### Notes

- Classification as opposed to Decision
- Loss function simply counts errors (misclassifications)
- We consider all errors equally painful!
- More examples during the lab ...
- The final result is not that surprising, is it?

## A special case - Bayesian *classification*

- ▶ Bayesian classification is a special case of statistical decision theory:
  - ▶ Attribute vector  $\vec{x} = (x_1, x_2, \dots)$ : pixels 1, 2, ...
  - ▶ **State set  $\mathcal{S}$  = decision set  $\mathcal{D} = \{0, 1, \dots, 9\}$ .**
  - ▶ **State = actual class, Decision = recognized class**
  - ▶ Loss function:

$$l(s, d) = \begin{cases} 0, & d = s \\ 1, & d \neq s \end{cases}$$

$$\delta^*(\vec{x}) = \arg \min_d \sum_s \underbrace{l(s, d)}_{0 \text{ if } d=s} P(s|\vec{x}) = \arg \min_d \sum_{s \neq d} P(s|\vec{x})$$

Obviously  $\sum_s P(s|\vec{x}) = 1$ , then:

$$P(d|\vec{x}) + \sum_{s \neq d} P(s|\vec{x}) = 1$$

Inserting into above:

$$\delta^*(\vec{x}) = \arg \min_d [1 - P(d|\vec{x})] = \arg \max_d P(d|\vec{x})$$

---

### Notes

- Classification as opposed to Decision
- Loss function simply counts errors (misclassifications)
- We consider all errors equally painful!
- More examples during the lab ...
- The final result is not that surprising, is it?



## A special case - Bayesian *classification*

- ▶ Bayesian classification is a special case of statistical decision theory:
  - ▶ Attribute vector  $\vec{x} = (x_1, x_2, \dots)$ : pixels 1, 2, ...
  - ▶ **State set  $\mathcal{S}$  = decision set  $\mathcal{D} = \{0, 1, \dots, 9\}$ .**
  - ▶ **State = actual class, Decision = recognized class**
  - ▶ Loss function:

$$l(s, d) = \begin{cases} 0, & d = s \\ 1, & d \neq s \end{cases}$$

$$\delta^*(\vec{x}) = \arg \min_d \sum_s \underbrace{l(s, d)}_{0 \text{ if } d=s} P(s|\vec{x}) = \arg \min_d \sum_{s \neq d} P(s|\vec{x})$$

Obviously  $\sum_s P(s|\vec{x}) = 1$ , then:

$$P(d|\vec{x}) + \sum_{s \neq d} P(s|\vec{x}) = 1$$

Inserting into above:

$$\delta^*(\vec{x}) = \arg \min_d [1 - P(d|\vec{x})] = \arg \max_d P(d|\vec{x})$$

22 / 24

---

### Notes

- Classification as opposed to Decision
- Loss function simply counts errors (misclassifications)
- We consider all errors equally painful!
- More examples during the lab ...
- The final result is not that surprising, is it?

## A special case - Bayesian *classification*

- ▶ Bayesian classification is a special case of statistical decision theory:
  - ▶ Attribute vector  $\vec{x} = (x_1, x_2, \dots)$ : pixels 1, 2, ...
  - ▶ **State set  $\mathcal{S}$  = decision set  $\mathcal{D} = \{0, 1, \dots, 9\}$ .**
  - ▶ **State = actual class, Decision = recognized class**
  - ▶ Loss function:

$$l(s, d) = \begin{cases} 0, & d = s \\ 1, & d \neq s \end{cases}$$

$$\delta^*(\vec{x}) = \arg \min_d \sum_s \underbrace{l(s, d)}_{0 \text{ if } d=s} P(s|\vec{x}) = \arg \min_d \sum_{s \neq d} P(s|\vec{x})$$

Obviously  $\sum_s P(s|\vec{x}) = 1$ , then:

$$P(d|\vec{x}) + \sum_{s \neq d} P(s|\vec{x}) = 1$$

Inserting into above:

$$\delta^*(\vec{x}) = \arg \min_d [1 - P(d|\vec{x})] = \arg \max_d P(d|\vec{x})$$

---

### Notes

- Classification as opposed to Decision
- Loss function simply counts errors (misclassifications)
- We consider all errors equally painful!
- More examples during the lab ...
- The final result is not that surprising, is it?

## A special case - Bayesian *classification*

- ▶ Bayesian classification is a special case of statistical decision theory:
  - ▶ Attribute vector  $\vec{x} = (x_1, x_2, \dots)$ : pixels 1, 2, ...
  - ▶ **State set  $\mathcal{S}$  = decision set  $\mathcal{D} = \{0, 1, \dots, 9\}$ .**
  - ▶ **State = actual class, Decision = recognized class**
  - ▶ Loss function:

$$l(s, d) = \begin{cases} 0, & d = s \\ 1, & d \neq s \end{cases}$$

$$\delta^*(\vec{x}) = \arg \min_d \sum_s \underbrace{l(s, d)}_{0 \text{ if } d=s} P(s|\vec{x}) = \arg \min_d \sum_{s \neq d} P(s|\vec{x})$$

Obviously  $\sum_s P(s|\vec{x}) = 1$ , then:

$$P(d|\vec{x}) + \sum_{s \neq d} P(s|\vec{x}) = 1$$

Inserting into above:

$$\delta^*(\vec{x}) = \arg \min_d [1 - P(d|\vec{x})] = \arg \max_d P(d|\vec{x})$$

22 / 24

### Notes

- Classification as opposed to Decision
- Loss function simply counts errors (misclassifications)
- We consider all errors equally painful!
- More examples during the lab ...
- The final result is not that surprising, is it?

# References I

Further reading: Chapter 13 and 14 of [6]. Books [1] and [2] are classical textbooks in the field of pattern recognition and machine learning. An interesting insights into how people think and interact with probabilities are presented in [4] (in Czech as [5])

[1] Christopher M. Bishop.

*Pattern Recognition and Machine Learning.*

Springer Science+Business Media, New York, NY, 2006.

PDF freely downloadable.

[2] Richard O. Duda, Peter E. Hart, and David G. Stork.

*Pattern Classification.*

John Wiley & Sons, 2nd edition, 2001.

[3] Zdeněk Kotek, Petr Vysoký, and Zdeněk Zdráhal.

*Kybernetika.*

SNTL, 1990.

# References II

- [4] Leonard Mlodinow.  
*The Drunkard's Walk. How Randomness Rules Our Lives.*  
Vintage Books, 2008.
- [5] Leonard Mlodinow.  
*Život je jen náhoda. Jak náhoda ovlivňuje naše životy.*  
Slovart, 2009.
- [6] Stuart Russell and Peter Norvig.  
*Artificial Intelligence: A Modern Approach.*  
Prentice Hall, 3rd edition, 2010.  
<http://aima.cs.berkeley.edu/>.