

## Exercise - combinatorics

- ▶ Assume a propositional logic over  $n$  variables. Compute the number of non-equivalent:
  - ▶ Monotone conjunctions
  - ▶ Conjunctions
- ▶ Can we similarly compute the number of non-equivalent  $s$ -DNFs? Can we put an upper bound on such number?

## Exercise - union bound (a.k.a. Boole's inequality)

- ▶  $P(A \cup B) \leq P(A) + P(B)$ 
  - ▶ Generalize to  $P(\bigcup_i A_i)$ .
  - ▶ How does it relate to the inclusion-exclusion principle?

## Exercise - Hoeffding's inequality

- ▶ Theorem: For random variables  $X_i, i = 1, \dots, n$  such that  $0 \leq X_i \leq 1$  and an  $\epsilon > 0$ , it holds that:  
$$P(\bar{X} - \mathbb{E}\bar{X} \geq \epsilon) \leq \exp(-2n\epsilon^2).$$
- ▶ Derive a similar bound for:  $P(|\bar{X} - \mathbb{E}\bar{X}| \geq \epsilon)$

## Exercise - Prosecutor's fallacy

- ▶ A DNA sample is found at a crime place.
- ▶ A match is found in a DB of 20000 people.
- ▶ The prosecutor explains that the probability that two profiles match by chance is only 1 in 10000.
- ▶ How strong evidence is it against the suspect? I.e. what is the probability of getting at least one match?