

3D Computer Vision

Radim Šára Martin Matoušek

Center for Machine Perception
Department of Cybernetics
Faculty of Electrical Engineering
Czech Technical University in Prague

<https://cw.fel.cvut.cz/wiki/courses/tdv/start>

<http://cmp.felk.cvut.cz>

<mailto:sara@cmp.felk.cvut.cz>

phone ext. 7203

rev. September 20, 2022



Open Informatics Master's Course

Perspective Camera

- 2.1 Basic Entities: Points, Lines
- 2.2 Homography: Mapping Acting on Points and Lines
- 2.3 Canonical Perspective Camera
- 2.4 Changing the Outer and Inner Reference Frames
- 2.5 Projection Matrix Decomposition
- 2.6 Anatomy of Linear Perspective Camera
- 2.7 Vanishing Points and Lines


covered by

[H&Z] Secs: 2.1, 2.2, 3.1, 6.1, 6.2, 8.6, 2.5, Example: 2.19

► Basic Geometric Entities, their Representation, and Notation

- entities have names and representations
- names and their components:

entity	in 2-space	in 3-space
point	$m = (u, v)$	$X = (x, y, z)$
line	n	O
plane		π, φ



- associated vector representations

$$\mathbf{m} = \begin{bmatrix} u \\ v \end{bmatrix} = [u, v]^T, \quad \mathbf{X} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad \mathbf{n}$$

will also be written in an 'in-line' form as $\mathbf{m} = (u, v)$, $\mathbf{X} = (x, y, z)$, etc.

- vectors are always meant to be columns $\mathbf{x} \in \mathbb{R}^{n \times 1}$
- associated homogeneous representations

$$\underline{\mathbf{m}} = [m_1, m_2, m_3]^T, \quad \underline{\mathbf{X}} = [x_1, x_2, x_3, x_4]^T, \quad \underline{\mathbf{n}}$$

'in-line' forms: $\underline{\mathbf{m}} = (m_1, m_2, m_3)$, $\underline{\mathbf{X}} = (x_1, x_2, x_3, x_4)$, etc.

- matrices are $\mathbf{Q} \in \mathbb{R}^{m \times n}$, linear map of a $\mathbb{R}^{n \times 1}$ vector is $\mathbf{y} = \mathbf{Q}\mathbf{x}$
- j -th element of vector \mathbf{m}_i is $(\mathbf{m}_i)_j$; element i, j of matrix \mathbf{P} is \mathbf{P}_{ij}

► Image Line (in 2D)


a finite line in the 2D (u, v) plane

$$(u, v) \in \mathbb{R}^2 \quad \text{s.t.} \quad a^2 + b^2 = 1 \quad a u + b v + c = 0$$

has a parameter (homogeneous) vector

$$\underline{\mathbf{n}} \simeq (a, b, c), \quad \|\underline{\mathbf{n}}\| \neq 0$$

and there is an equivalence class for $\lambda \in \mathbb{R}, \lambda \neq 0$ $(\lambda a, \lambda b, \lambda c) \simeq (a, b, c)$



'Finite' lines

- standard representative for finite $\underline{\mathbf{n}} = (n_1, n_2, n_3)$ is $\lambda \underline{\mathbf{n}}$, where $\lambda = \frac{1}{\sqrt{n_1^2 + n_2^2}}$
assuming $n_1^2 + n_2^2 \neq 0$; $\mathbf{1}$ is the unit, usually $\mathbf{1} = 1$

'Infinite' line

- we augment the set of lines for a special entity called the **line at infinity** (ideal line)

$$\underline{\mathbf{n}}_\infty \simeq (0, 0, 1) \quad (\text{standard representative})$$


- the set of equivalence classes of vectors in $\mathbb{R}^3 \setminus (0, 0, 0)$ forms the projective space \mathbb{P}^2 a set of rays $\rightarrow 21$
- line at infinity is a proper member of \mathbb{P}^2
- I may sometimes wrongly use $=$ instead of \simeq , if you are in doubt, ask me

► Image Point

Finite point $\underline{\mathbf{m}} = (u, v)$ is incident on a finite line $\underline{\mathbf{n}} = (a, b, c)$ iff

$$a u + b v + c \neq 0$$

iff = works either way!



can be rewritten as (with scalar product): $(u, v, \mathbf{1}) \cdot (a, b, c) = \underline{\mathbf{m}}^\top \underline{\mathbf{n}} = 0$

'Finite' points

- a finite point is also represented by a homogeneous vector $\underline{\mathbf{m}} \simeq (u, v, \mathbf{1})$, $\|\underline{\mathbf{m}}\| \neq 0$
- the equivalence class for $\lambda \in \mathbb{R}, \lambda \neq 0$ is $(m_1, m_2, m_3) = \lambda \underline{\mathbf{m}} \simeq \underline{\mathbf{m}}$
- the standard representative for finite point $\underline{\mathbf{m}}$ is $\lambda \underline{\mathbf{m}}$, where $\lambda = \frac{1}{m_3}$
- when $\mathbf{1} = 1$ then units are pixels and $\lambda \underline{\mathbf{m}} = (u, v, 1)$
- when $\mathbf{1} = f$ then all elements have a similar magnitude, $f \sim$ image diagonal

assuming $m_3 \neq 0$

use $\mathbf{1} = 1$ unless you know what you are doing;

all entities participating in a formula must be expressed in the same units

'Infinite' points


- we augment for **points at infinity** (ideal points) $\underline{\mathbf{m}}_\infty \simeq (m_1, m_2, 0)$
- all such points lie on the line at infinity (ideal line) $\underline{\mathbf{n}}_\infty \simeq (0, 0, 1)$, i.e. $\underline{\mathbf{m}}_\infty^\top \underline{\mathbf{n}}_\infty = 0$

proper members of \mathbb{P}^2


► Line Intersection and Point Join

The point of **intersection** m of image lines n and n' , $n \neq n'$ is

$$\underline{\mathbf{m}} \simeq \underline{\mathbf{n}} \times \underline{\mathbf{n}'}$$



proof: If $\underline{\mathbf{m}} = \underline{\mathbf{n}} \times \underline{\mathbf{n}'}$ is the intersection point, it must be incident on both lines. Indeed, using known equivalences from vector algebra

$$\begin{aligned} \underline{\mathbf{n}}^T (\underline{\mathbf{n}} \times \underline{\mathbf{n}'}) &\equiv \underline{\mathbf{n}'^T} (\underline{\mathbf{n}} \times \underline{\mathbf{n}'}) \equiv 0 \\ \underbrace{\underline{\mathbf{n}}^T}_{\underline{\mathbf{m}}} \underbrace{(\underline{\mathbf{n}} \times \underline{\mathbf{n}'})}_{\underline{\mathbf{m}}} &= \underbrace{\underline{\mathbf{n}'^T}}_{\underline{\mathbf{m}}} \underbrace{(\underline{\mathbf{n}} \times \underline{\mathbf{n}'})}_{\underline{\mathbf{m}}} = 0 \\ \underline{\mathbf{n}}^T \cdot \underline{\mathbf{m}} &= 0 \end{aligned}$$
A diagram showing a line labeled n passing through two points, m and m' .

The **join** n of two image points m and m' , $m \neq m'$ is

$$\underline{\mathbf{n}} \simeq \underline{\mathbf{m}} \times \underline{\mathbf{m}'}$$

Parallel lines intersect (somewhere) on the line at infinity $\underline{\mathbf{n}}_\infty \simeq (0, 0, 1)$:

$$a u + b v + c = 0,$$


$$a u + b v + d = 0,$$

$$d \neq c$$

$$(a, b, c) \times (a, b, d) \simeq (b, -a, 0)$$

- all such intersections lie on $\underline{\mathbf{n}}_\infty$
- line at infinity therefore represents the set of (unoriented) directions in the plane
- Matlab: `m = cross(n, n_prime);`

► Homography in \mathbb{P}^2




Projective plane \mathbb{P}^2 : Vector space of dimension 3 excluding the zero vector, $\mathbb{R}^3 \setminus (0,0,0)$, factorized to linear equivalence classes ('rays'), $\underline{x} \simeq \lambda \underline{x}$, $\lambda \neq 0$ including 'points at infinity'

we call $\underline{x} \in \mathbb{P}^2$ 'points'

Homography in \mathbb{P}^2 : Non-singular linear mapping in \mathbb{P}^2

$$\underline{x}' \simeq \mathbf{H} \underline{x}, \quad \mathbf{H} \in \mathbb{R}^{3,3} \text{ non-singular}$$

an analogic definition for \mathbb{P}^3



lines of points are mapped to lines of points

concurrent = intersecting at a point

e.g. line intersection points mapped to line intersection points

$$H \text{ pts} \leftrightarrow H^T \sim H \text{ line}$$


$$\mathbf{H} \in \text{SL}(3)$$

Defining properties

- collinear points are mapped to collinear points
- concurrent lines are mapped to concurrent lines
- and point-line incidence is preserved

- \mathbf{H} is a 3×3 non-singular matrix, $\lambda \mathbf{H} \simeq \mathbf{H}$ equivalence class, 8 degrees of freedom
- homogeneous matrix representative: $\det \mathbf{H} = 1$
- what we call homography here is often called 'projective collineation' in mathematics

► Mapping 2D Points and Lines by Homography



$$\underline{\mathbf{m}}' \simeq \mathbf{H} \underline{\mathbf{m}} \quad (\text{image) point}$$

$$\underline{\mathbf{n}}' \simeq \mathbf{H}^{-\top} \underline{\mathbf{n}} \quad (\text{image) line}$$

$$\mathbf{H}^{-\top} = (\mathbf{H}^{-1})^\top = (\mathbf{H}^\top)^{-1}$$

- incidence is preserved. $(\underline{\mathbf{m}}')^\top \underline{\mathbf{n}}' \simeq \underline{\mathbf{m}}^\top \mathbf{H}^\top (\mathbf{H}^{-\top} \underline{\mathbf{n}}) = \underline{\mathbf{m}}^\top \underline{\mathbf{n}} = 0$

$$\mathbf{H}^\top \mathbf{M} = \mathbf{I} \cdot \lambda$$

$$\mathbf{H}^\top \mathbf{H} \simeq \mathbf{I}$$

Mapping a finite 2D point $\mathbf{m} = (u, v)$ to $\underline{\mathbf{m}} = (u', v')$

- extend the Cartesian (pixel) coordinates to homogeneous coordinates, $\underline{\mathbf{m}} = (u, v, 1)$
- map by homography, $\underline{\mathbf{m}}' = \mathbf{H} \underline{\mathbf{m}}$
- if $m'_3 \neq 0$ convert the result $\underline{\mathbf{m}}' = (m'_1, m'_2, m'_3)$ back to Cartesian coordinates (pixels),

$$u' = \frac{m'_1}{m'_3} \mathbf{1}, \quad v' = \frac{m'_2}{m'_3} \mathbf{1}$$

- note that, typically, $m'_3 \neq 1$
- an infinite point $\underline{\mathbf{m}} = (u, v, 0)$ maps the same way



$m'_3 = 1$ when \mathbf{H} is affine

Thank You