

# Homework (A8B17CAS)

## Problem Set 3

December 12, 2023

### 1 Assignment

P3-A Define MATLAB function `rot90` that takes a matrix (a list of lists) and rotates it counter-clockwise. *E.g.* an original matrix, `matA`, the output/transformed matrix, `matB = rot90(matA)`:

$$\text{matA} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}, \quad \text{matB} = \begin{bmatrix} 3 & 6 \\ 2 & 5 \\ 1 & 4 \end{bmatrix}. \quad (1)$$

Task: Write your definition of `rot90` to your notebook.

A hint: Take a look at MATHEMATICA command `Reverse`.

**(1 points)**

P3-B Assume a function  $f(x) = x^5 e^{-x}$ . Compute its first 4 derivatives. The function and their derivative are in Figure 1. Get the  $x$ -coordinates of the 5 shown points. You can use the command `FindRoot`. It requires a starting point. Use the starting points from the list/vector: `{5.1, 3, 1.5, 1, 0.5}`;




Figure 1: Function  $f(x)$  and its first 4 derivatives.

Task: Get a list of  $x$ -coordinates of the 5 shown points. (Transform a list of rules that you get from `FindRoot` to a list of numbers).

A hint: Additional command that can be useful is `MapThread`, and maybe `NestList`.

**(2 points)**

P3-C Define a function `myMat [p-]` that for a given  $p \geq 3$ ,  $p \in \mathbb{Z}$  generates the following square matrix:

$$\begin{bmatrix} -p & 0 & \dots & 0 & & & \\ \vdots & \vdots & \ddots & \vdots & & & \\ \vdots & 0 & \dots & 0 & & & \\ -1 & 1 & \dots & 1 & 1 & \dots & 1 \\ 0 & 1 & \dots & 1 & 1 & \dots & 1 \end{bmatrix}. \quad (2)$$

E.g.

$$\text{myMat}[5] = \begin{bmatrix} -5 & 0 & 0 & 0 & 12 & 10 \\ -4 & 0 & 0 & 0 & 20 & 18 \\ -3 & 0 & 0 & 0 & 20 & 14 \\ -2 & 0 & 0 & 0 & 18 & 20 \\ -1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}, \text{myMat}[7] = \begin{bmatrix} -7 & 0 & 0 & 0 & 18 & 20 & 14 & 12 \\ -6 & 0 & 0 & 0 & 18 & 13 & 11 & 11 \\ -5 & 0 & 0 & 0 & 19 & 15 & 14 & 13 \\ -4 & 0 & 0 & 0 & 16 & 10 & 12 & 20 \\ -3 & 0 & 0 & 0 & 10 & 15 & 16 & 10 \\ -2 & 0 & 0 & 0 & 15 & 20 & 11 & 10 \\ -1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}. \quad (3)$$

Where  $\mathbf{R} \in \mathbb{R}^{(p-1) \times (p-3)}$  is a matrix of random integers from a uniform distribution on a set  $\{10, \dots, 20\}$ . Matrix of zeros,  $\mathbf{O} \in \mathbb{R}^{(p-1) \times 3}$  and matrix of ones  $\mathbf{1} \in \mathbb{R}^{2 \times p}$ .

Task: Write your definition of `myMat` to your notebook.

**(2 points)**

P3-D Assume  $n \in \mathbb{N}$ . Generate a list of  $n$  2D points (a matrix with 2 columns and  $n$  rows). Denote the list as `ptsRhoPhi`, it is a list of the form  $\{\{\rho_1, \varphi_1\}, \{\rho_2, \varphi_2\}, \dots, \{\rho_n, \varphi_n\}\}$ . Generate  $\rho_i$ s as a random variable from the [normal distribution](#)  $\mathcal{N}(10, 1)$  – `RandomVariate[NormalDistribution[10, 1]]` and  $\varphi_i$ s as a random variable from the [continuous uniform distribution](#)  $\mathcal{U}(0, 2\pi)$  – `RandomVariate[UniformDistribution[0, 2 Pi]]`.

Create matrices/lists `ptsXY` and `ptsXYSorted` such that:

- the list `ptsXY` contains pairs  $\{x_i, y_i\}$ , where  $x_i = \rho_i \cos(\varphi_i)$  and  $y_i = \rho_i \sin(\varphi_i)$ .
- the list `ptsXYSorted` is sorted `ptsXY` by the angle of the point in plain.

Task: For a given  $n$ , create list `ptsRhoPhi`. Then create lists `ptsXY` and `ptsXYSorted`. Write the procedure to your notebook.

A hint: You can use e.g. [ArcTan](#) to get the angle from  $x, y$  coordinates.

You can plot the resulting vectors using:

```
ListPlot[{ptsXY, ptsXYSorted}, Joined -> True, AspectRatio -> 1, AxesLabel -> {"x", "y"}, PlotLegends -> {"ptsXY", "ptsXYSorted"}]
```

Figure 2 shows an example of such a plot.




Figure 2: A plot of `ptsXY` and `ptsXYSorted` for  $n = 30$ .

**(2 points)**

P3-E Generate Pascal's triangle and its modified version. Do not use the [Binomial](#) command. Compute the rows iteratively so that your function takes one row (a list of numbers) and returns the next row of numbers. Use your function with the command [NestList](#) to get a list of the first few rows.

Define a second function: instead of the sum  $x_1 + x_2$  of the above numbers  $x_1, x_2$  returns  $x_1 + 2x_2$ . Use the function with command `NestList` to get the first few rows of such a triangle.

You can use the postfix command `// Column[#, Center] &` to show the list of rows in a triangle form.




Figure 3: The Pascal's triangle and its modified version.

Task: Compute the first few rows of the ordinary Pascal's triangle and its modified version.

A hint: Useful commands to look at are `Partition` and `Join`.

**(3 points)**

## 2 Instructions

Complete all the assignments till

- January 7 2024, 23:59.

Write your solutions into a notebook called `Problem3_yourusername.nb` and send it to [lukacjo1@fel.cvut.cz](mailto:lukacjo1@fel.cvut.cz) with subject "CAS:HW3". All the problems shall be solved by the students individually. Do not use functions from MATHEMATICA Packages.

Contact us at [lukacjo1@fel.cvut.cz](mailto:lukacjo1@fel.cvut.cz) with any questions. The team of teachers wishes you good luck in solving this.