

**First homework assignment** announced 7. 3. 2025, due 31. 3. 2025

**Q1:** Consider the 2-qubit state  $|\psi^+\rangle := \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$ . Calculate the expectation values in this state for the operators

a)  $H \otimes H$ , and **(1.5p)**

b)  $H \otimes \sigma_z$ . **(1.5p)**

Here  $H$  is the Hadamard operator,  $H := \frac{1}{\sqrt{2}}(\sigma_x + \sigma_z)$ , and  $\sigma_x, \sigma_z$  the ordinary Pauli operators.

**Q2:** Consider the Hamiltonian operator of a 2-dimensional quantum harmonic oscillator

$$H = \frac{1}{2m}(p_x^2 + p_y^2) + \frac{1}{2}m\omega^2(x^2 + y^2).$$

This can be written as the sum of the Hamiltonian of two one-dimensional oscillators

$$H = H_x + H_y, \quad H_j = \frac{1}{2m}p_j^2 + \frac{1}{2}m\omega^2 j^2,$$

for  $j = x, y$ . The momentum and position operators satisfy the commutation relations:  $[x, p_x] = [y, p_y] = i\hbar$ , while the rest are zero, i.e.  $[x, y] = [x, p_y] = [y, p_x] = [p_x, p_y] = 0$ .

a) Does  $H_x$  and  $H_y$  commute? **(1.5p)**

b) Can you construct a non-trivial (i.e. not zero, the identity or a factor of  $H$  itself) operator, using only polynomial combinations of  $x, y, p_x$  and  $p_y$ , that commutes with the full Hamiltonian  $H$ ? If possible, what does this tell you about this quantity? **(1.5p)**

**Q3:** Alice and Bob are studying a 3-dimensional quantum system  $|\psi\rangle \in \mathbb{C}^3$ . Alice measures an observable that can take values red, green and blue (or  $r, g$  and  $b$ ) while Bob measures an observable that gives values sweet, tangy or umami (or  $s, t$  and  $u$ ). If Alice find the result  $r$ , then Bob finds that he finds the corresponding states  $s, t$  or  $u$  with probabilities  $0, p$  and  $1 - p$ , respectively. If on the other hand, Alice finds  $g$ , Bob's probabilities becomes  $q, 0$  and  $1 - q$ , for the values  $s, t$ , and  $u$ , respectively.

a) Which combinations of values are allowed for  $p$  and  $q$ ? **(2p)**

b) What are Bob's probabilities if Alice finds the result  $b$ ? **(2p)**