

# Quantum Computing - Exercise Sheet 7

## Quantum Random Walks

1. At each time step, a quantum walk corresponds to a unitary map  $U \in U(N)$  such that

$$U : \mathcal{H}_G \rightarrow \mathcal{H}_G$$

$$|x\rangle \mapsto a|x-1\rangle + b|x\rangle + c|x+1\rangle$$

Show that  $U$  is unitary if and only if one of the following three conditions is true: (a)  $|a| = 1, b = c = 0$ , (b)  $|b| = 1, a = c = 0$ , (c)  $|c| = 1, a = b = 0$ .

2. Demonstrate that the shift operator  $S$ , as defined as

$$S = (|0\rangle\langle 0| \otimes \sum_{x=-\infty}^{+\infty} |x+1\rangle\langle x|) + (|1\rangle\langle 1| \otimes \sum_{x=-\infty}^{+\infty} |x-1\rangle\langle x|)$$

is equivalent to

$$S|i, x\rangle = \begin{cases} |0, x+1\rangle & \text{if } i = 0, \\ |1, x-1\rangle & \text{if } i = 1. \end{cases}$$

3. Consider a one-dimensional quantum walk on  $\mathbb{Z}$  where the coin operator  $C$  is parameterized by an angle  $\theta$  as :

$$c(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$$

The walker starts at  $x = 0$  with initial state  $|\psi_0\rangle = |i\rangle \otimes |x=0\rangle$  and  $|i\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ . Use the shift operator as defined before a) Calculate the state of the system  $|\psi_1\rangle$  after 1 step.

b) Compute the probabilities  $p(x=1)$  and  $p(x=-1)$  that the walker is at  $x=1$  or  $x=-1$ .

c) Determine for which values of  $\theta$  (if they exist) the quantum walks are unbiased ( $p(x=1) = p(x=-1)$ )

d) Prove whether the Hadamard walker is biased or not.

4. Starting at the state  $|\psi_0\rangle = |0\rangle|0\rangle$ , obtain the successive states up  $|\psi_4\rangle$  for the Hadamard walker on the finite subset of  $\mathbb{Z}$ .

5. Consider

$$H_G^{(2)} = \sum_{\omega=1}^2 \sum_{(i,j) \in E(G)} (|i\rangle\langle j|_{\omega} + |j\rangle\langle i|_w)$$

where  $E(G) = \{(1,2) \text{ and } (2,1)\}$ . This is the Hamiltonian for 2 particles on this  $G$ .

a) Assume we have distinguishable walkers. Compute the evolution of the initial state  $|\psi_0\rangle = |1,2\rangle$  under the Hamiltonian  $H_G^{(2)}$ .

b) Assuming the walkers are distinguishable, now compute the evolution for the fermionic state  $|\psi_0\rangle = \frac{1}{\sqrt{2}}(|1,2\rangle - |2,1\rangle)$