

# AE4M33RZN, Fuzzy logic: Fuzzy relations

Radomír Černoč

radomir.cernoch@fel.cvut.cz

26/11/2012

Faculty of Electrical Engineering, CTU in Prague

# Plan of the lecture

## Properties of fuzzy sets

- Fuzzy implication and fuzzy properties

- Fuzzy set inclusion and crisp predicates

## Intermission: Probabilistic vs. fuzzy

## Binary fuzzy relations

- Quick revision of crisp relations

- Fuzzyfication of crisp relations

- Projection and cylindrical extension

- Composition of fuzzy relations

- Properties of fuzzy relations

- Properties of fuzzy composition

## Extensions

## Biblography

## Organizational:

- Next week, there will be a short test (max 5 points).
- This week we are having the last **theoretical lecture**.

# Fuzzy implication

We already know *fuzzy negation*  $\neg_{\circ}$ , *fuzzy conjunction*  $\wedge_{\circ}$  and *fuzzy disjunction*  $\vee_{\circ}$ . What about other operators?

# Fuzzy implication

We already know *fuzzy negation*  $\overset{\circ}{\neg}$ , *fuzzy conjunction*  $\overset{\circ}{\wedge}$  and *fuzzy disjunction*  $\overset{\circ}{\vee}$ . What about other operators?

## Definition

*Fuzzy implication* is any function

$$\overset{\circ}{\Rightarrow} : [0, 1]^2 \rightarrow [0, 1] \quad (1)$$

which overlaps with the boolean implication on  $x, y \in \{0, 1\}$ :

$$(x \overset{\circ}{\Rightarrow} y) = (x \Rightarrow y) . \quad (2)$$

# Residue implication

Despite the lack of a uniform definition of fuzzy implication, there is a useful class of implications:

## Defintion



The *R-implication* (residuum, „*reziduovaná implikace*“) is a function obtained from a fuzzy T-norm:

$$\alpha \overset{R}{\underset{\circ}{\Rightarrow}} \beta = \sup\{\gamma \mid \alpha \underset{\circ}{\wedge} \gamma \leq \beta\} \quad (\text{RI})$$

# R-implication: Examples (1)

*Standard implication* (Gödel) is derived from (RI) using the standard conjunction  $\wedge_S$ :



$$\alpha \xrightarrow[S]{R} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ \beta & \text{otherwise} \end{cases} \quad (3)$$



## R-implication: Examples (2)

*Lukasiewicz implication* is derived from (RI) using the Łukasiewicz conjunction  $\wedge_L$ :

$$\alpha \xrightarrow[\text{L}]{\text{R}} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ 1 - \alpha + \beta & \text{otherwise} \end{cases} \quad (4)$$







## R-implication: Examples (3)

*Algebraic implication* (Gougen, Gaines) is derived from (RI) using the algebraic conjunction  $\wedge_A$ :

$$\alpha \xrightarrow[A]{R} \beta = \begin{cases} 1 & \text{if } \alpha \leq \beta \\ \frac{\beta}{\alpha} & \text{otherwise} \end{cases} \quad (5)$$



# R-implication: Properties

## Theorem 209.

Let  $\triangleleft_{\circ}$  be a continuous fuzzy conjunction. Then R-implication satisfies:

$$\alpha \xrightarrow[\circ]{R} \beta = \mathbf{1} \text{ iff } \alpha \leq \beta \quad (I1)$$

$$\mathbf{1} \xrightarrow[\circ]{R} \beta = \beta \quad (I2)$$

$$\alpha \xrightarrow[\circ]{R} \beta \text{ is not increasing in } \alpha \text{ and not decreasing in } \beta \quad (I3)$$

# R-implication: Properties

## Proof of theorem 209.

Let's denote  $\{\gamma \mid \alpha \underset{\circ}{\wedge} \gamma \leq \beta\} = \gamma$ .

- Proving (I3) uses monotonicity: Increasing  $\alpha$  can only shrink  $\gamma$  and increasing  $\beta$  can only enlarge  $\gamma$ .
- Proving (I2) is easy:  $1 \xrightarrow{\underset{\circ}{R}} \beta = \sup\{\gamma \mid 1 \underset{\circ}{\wedge} \gamma \leq \beta\}$ . From definition of  $\underset{\circ}{\wedge}$ , we write  $1 \xrightarrow{\underset{\circ}{R}} \beta = \sup\{\gamma \mid \gamma \leq \beta\} = \beta$ .

## Proof of theorem 209 (contd.).

- For (I1) one needs to check 2 cases:
  - If  $\alpha \leq \beta$ , then  $1 \in \gamma$ , because  $\alpha \wedge_{\circ} 1 = \alpha \leq \beta$  and therefore the condition  $\alpha \wedge_{\circ} \gamma \leq \beta$  is true for all possible values of  $\gamma$ .
  - If  $\alpha > \beta$ , then  $1 \notin \gamma$ , because  $\alpha \wedge_{\circ} 1 = \alpha > \beta$  and therefore the condition  $\alpha \wedge_{\circ} \gamma \leq \beta$  is false for  $\gamma = 1$ .

# S-implication

## Defintion

The *S-implication* is a function obtained from a fuzzy disjunction  $\overset{\circ}{\vee}$ :

$$\alpha \underset{\circ}{\overset{\text{S}}{\Rightarrow}} \beta = \underset{\text{S}}{\neg} \alpha \overset{\circ}{\vee} \beta \quad (\text{SI})$$

# S-implication

## Definition

The *S-implication* is a function obtained from a fuzzy disjunction  $\overset{\circ}{\vee}$ :

$$\alpha \overset{\text{S}}{\underset{\circ}{\Rightarrow}} \beta = \underset{\text{S}}{\neg} \alpha \overset{\circ}{\vee} \beta \quad (\text{SI})$$

## Example

*Kleene-Dienes* implication from  $\overset{\text{S}}{\underset{\circ}{\vee}}$

$$\alpha \overset{\text{S}}{\underset{\circ}{\Rightarrow}} \beta = \max(1 - \alpha, \beta) \quad (6)$$

# Generalized fuzzy inclusion

# Generalized fuzzy inclusion

Previously, we used the logical negation  $\neg$  to define the set complement, the conjunction  $\wedge$  to define the set intersection, etc.

Can we use the implication  $\Rightarrow$  to define the the fuzzy inclusion?



# Generalized fuzzy inclusion

Previously, we used the logical negation  $\neg$  to define the set complement, the conjunction  $\wedge$  to define the set intersection, etc.

Can we use the implication  $\Rightarrow$  to define the the fuzzy inclusion?

## Definition

The *generalized fuzzy inclusion*  $\underset{\circ}{\subseteq}$  is a function that assigns a degree to the the inclusion of set  $A \in \mathbb{F}(\Delta)$  in set  $B \in \mathbb{F}(\Delta)$ :

$$A \underset{\circ}{\subseteq} B = \inf\{A(\mathbf{x}) \underset{\circ}{\Rightarrow} B(\mathbf{x}) \mid \mathbf{x} \in \Delta\} \quad (7)$$

# Generalized fuzzy inclusion: Example

# Fuzzy inclusion (non-generalized)

## Definition

The *fuzzy inclusion*  $\subseteq$  is a predicate (assigns a true/false value) which hold for two fuzzy sets  $A, B \in \mathbb{F}(\Delta)$  iff

$$\mu_A(\mathbf{x}) \leq \mu_B(\mathbf{x}) \text{ for all } \mathbf{x} \in \Delta. \quad (8)$$

# Fuzzy inclusion (non-generalized)

In vertical representation, the definition has a straightforward equivalent:

$$\mu_A \leq \mu_B \quad (9)$$

# Fuzzy inclusion (non-generalized)

In vertical representation, the definition has a straightforward equivalent:

$$\mu_A \leq \mu_B \quad (9)$$

In horizontal representation, there is a theorem:

## Theorem 219.

Let  $A, B \in \mathbb{F}(\Delta)$  if and only if

$$R_A(\alpha) \subseteq R_B(\alpha) \text{ for all } \alpha \in [0, 1] . \quad (10)$$

# Fuzzy inclusion (non-generalized)

## Proof of theorem 219.

- $\Rightarrow$  Assume  $A \subseteq B$  and  $\mathbf{x} \in R_A(\alpha)$  for some value  $\alpha$ . If  $\alpha \leq A(\mathbf{x})$ , then  $A(\mathbf{x}) \leq B(\mathbf{x})$  (from the definition of  $A \subseteq B$ ) and therefore  $\mathbf{x} \in R_B(\alpha)$  and  $R_A(\alpha) \subseteq R_B(\alpha)$ .
- $\Leftarrow$  Assume  $R_A(\alpha) \subseteq R_B(\alpha)$ . Firstly recall the horizontal-vertical translation formula:  $\mu_A(\mathbf{x}) = \sup\{\alpha \in [0,1] \mid \mathbf{x} \in R_A(\alpha)\}$ . Since  $\{\alpha \mid \mathbf{x} \in R_A(\mathbf{x})\} \subseteq \{\alpha \mid \mathbf{x} \in R_B(\mathbf{x})\}$ , the inequality  $A(\mathbf{x}) \leq \sup\{\alpha \mid \mathbf{x} \in R_B(\mathbf{x})\} \leq B(\mathbf{x})$  holds.

# Cutworthiness

We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

## Cutworthiness

Let  $P$  be a predicate (returns true/false) over fuzzy sets.  $P$  is called *cutworthy* („řezově dědičná vlastnost“) if the implication holds:

$$P(A_1, \dots, A_n) \Rightarrow P(R_{A_1}(\alpha), \dots, R_{A_n}(\alpha)) \text{ for all } \alpha \in [0, 1] \quad (11)$$



We ended up with 2 equal definitions of set inclusion: using vertical and horizontal representation. Can we generalize this?

## Cutworthiness

Let  $P$  be a predicate (returns true/false) over fuzzy sets.  $P$  is called *cutworthy* („řezově dědičná vlastnost“) if the implication holds:

$$P(A_1, \dots, A_n) \Rightarrow P(R_{A_1}(\alpha), \dots, R_{A_n}(\alpha)) \text{ for all } \alpha \in [0, 1] \quad (11)$$

There is a related notion: We define  $P$  as *cut-consistent* („řezově konzistentní“) using the same definition, but replacing  $\Rightarrow$  with  $\Leftrightarrow$ .

# Cutworthiness: Examples

- The theorem 219 can be stated as: “Set inclusion is cut-consistent.”

## Brain teasers

- *Strong normality* of  $A$  is defined as  $A(x) = 1$  for some  $x \in \Delta$ .  
????
- *Being crisp* is  
????


# Cutworthiness: Examples

- The theorem 219 can be stated as: “Set inclusion is cut-consistent.”

## Brain teasers



- *Strong normality* of  $A$  is defined as  $A(\mathbf{x}) = 1$  for some  $\mathbf{x} \in \Delta$ .  
Strong normality is **cut-consistent**:  $A$  is strongly-normal iff every its cut is non-empty iff every cut strongly normal.
- *Being crisp* is **cutworthy, but not cut-consistent**: Every cut is crisp by definition, therefore cutworthiness. But even **non-crisp sets** have crisp cuts, therefore the property is not not cut-consistent.

# Google: “fuzzy”



Sources: M. Taylor's Weblog, M. Taylor's Weblog, Eddie's Trick Shop.

# Google: “probability”



Sources: Life123, WhatWeKnowSoFar, Probability Problems.

# Fuzzy vs. probability

- *Vagueness vs. uncertainty.*

# Fuzzy vs. probability

- *Vagueness vs. uncertainty.*
- Fuzzy logic is *functional*.

## Definition

A *binary crisp relation*  $R$  from  $X$  onto  $Y$  is a subset of the cartesian product  $X \times Y$ :

$$R \in \mathbb{P}(X \times Y) \quad (12)$$



## Definition

A *binary crisp relation*  $R$  from  $X$  onto  $Y$  is a subset of the cartesian product  $X \times Y$ :

$$R \in \mathbb{P}(X \times Y) \quad (12)$$

## Definition

The *inverse relation*  $R^{-1}$  to  $R$  is a relation from  $Y$  to  $X$  s.t.

$$R^{-1} = \{(y, x) \in Y \times X \mid (x, y) \in R\} \quad (13)$$

# Crisp relations: Inverse

## Definition

Let  $X, Y, Z$  be sets. Then the *compound* of relations  $R \subseteq X \times Y, S \subseteq Y \times Z$  is the relation

$$R \circ S = \{(x, z) \in X \times Z \mid (x, y) \in R \text{ and } (y, z) \in S \text{ for some } y\} \quad (14)$$

## Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property

using logical connectives

using set axioms

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$



# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$
transitive	$(x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$	$R \circ R \subseteq R$

# Crisp relations: Properties

The *equality* relation on  $\Delta$  is  $E = \{(x, x) \mid x \in \Delta\}$ .

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using logical connectives	using set axioms
reflexive	$\forall x. (x, x) \in R$	$E \subseteq R$
symmetric	$(x, y) \in R \Rightarrow (y, x) \in R$	$R = R^{-1}$
anti-symmetric	$(x, y) \in R \wedge (y, z) \in R \Rightarrow y = z$	$R \cap R^{-1} \subseteq E$
transitive	$(x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$	$R \circ R \subseteq R$
partial order	reflexive, transitive and anti-symmetric	
equivalence	reflexive, transitive and symmetric	

## Definition

A *binary fuzzy relation*  $R$  from  $X$  onto  $Y$  is a fuzzy subset on the universe  $X \times Y$ .

$$R \in \mathbb{F}(X \times Y) \quad (15)$$

## Definition

The *fuzzy inverse* relation  $R^{-1} \in \mathbb{F}(Y \times X)$  to  $R \in \mathbb{F}(X \times Y)$ , s.t.

$$R(\mathbf{y}, \mathbf{x}) = R^{-1}(\mathbf{x}, \mathbf{y}) \quad (16)$$

## Defintion

Let  $R \in \mathbb{F}(X \times Y)$  be a fuzzy binary relation. The *first* and second projection of  $R$  is

$$R^{(1)}(\mathbf{x}) = \bigvee_{\mathbf{y} \in Y}^S R(\mathbf{x}, \mathbf{y}) \quad (17)$$

$$R^{(2)}(\mathbf{y}) = \bigvee_{\mathbf{x} \in X}^S R(\mathbf{x}, \mathbf{y}) \quad (18)$$

# Projection: Example

$R$	$y_1$	$y_2$	$y_3$	$y_4$	$y_5$	$y_6$	$R^{(1)}(x)$
$x_1$	0.1	0.2	0.4	0.8	1	0.8	?
$x_2$	0.2	0.4	0.8	1	0.8	0.6	?
$x_3$	0.4	0.8	1	0.8	0.4	0.2	?
$R^{(2)}(y)$	?	?	?	?	?	?	

# Projection: Example

$R$	$y_1$	$y_2$	$y_3$	$y_4$	$y_5$	$y_6$	$R^{(1)}(x)$
$x_1$	0.1	0.2	0.4	0.8	1	0.8	1
$x_2$	0.2	0.4	0.8	1	0.8	0.6	1
$x_3$	0.4	0.8	1	0.8	0.4	0.2	1
$R^{(2)}(y)$	0.4	0.8	1	0.8	0.4	0.2	

Sometimes there is a *total projection* defined as

$R^{(T)} = \bigvee_{x \in X} \bigvee_{y \in Y} R(x, y)$ . But we already know this notion as  $\text{Height}(R)$ .

# Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

# Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

## Definition

Let  $A \in \mathbb{F}(X)$  and  $B \in \mathbb{F}(Y)$  be fuzzy sets. The *cylindrical extension* („cylindrické rozšíření“, „kartézský součin fuzzy množin“) is defined as

$$A \times B(x, y) = A(x) \underset{S}{\wedge} B(y) \quad (19)$$



# Cylindrical extension

Can we reconstruct a fuzzy relation from its projections? There is an unique largest relation with prescribed projections:

## Definition

Let  $A \in \mathbb{F}(X)$  and  $B \in \mathbb{F}(Y)$  be fuzzy sets. The *cylindrical extension* („cylindrické rozšíření“, „kartézský součin fuzzy množin“) is defined as

$$A \times B(x, y) = A(x) \underset{S}{\wedge} B(y) \quad (19)$$

## Brain teaser

Why can't there be a relation  $Q$  bigger than  $A \times B$ , whose projections are  $Q^{(1)} = A$  and  $Q^{(2)} = B$ ?

## Cylindrical extension: Drawing

$$A(x) = \begin{cases} x - 1 & x \in [1, 2] \\ 3 - x & x \in [2, 3] \\ 0 & \text{otherwise} \end{cases}$$

$$B(x) = \begin{cases} x - 3 & x \in [3, 4] \\ 5 - x & x \in [4, 5] \\ 0 & \text{otherwise} \end{cases}$$

# Composition of fuzzy relations

## Definition

Let  $X, Y, Z$  be crisp sets.  $R \in \mathbb{F}(X \times Y)$ ,  $S \in \mathbb{F}(Y \times Z)$  and  $\wedge$  some fuzzy conjunction. Then the  $\circ$ -composition („ $\circ$ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge S(y, z) \quad (20)$$

# Composition of fuzzy relations

## Definition

Let  $X, Y, Z$  be crisp sets.  $R \in \mathbb{F}(X \times Y)$ ,  $S \in \mathbb{F}(Y \times Z)$  and  $\wedge$  some fuzzy conjunction. Then the  $\circ$ -composition („ $\circ$ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge S(y, z) \quad (20)$$

1. For infinite domains,  $\bigvee^S$  is computed using the **sup** instead of **max**.

# Composition of fuzzy relations

## Definition

Let  $X, Y, Z$  be crisp sets.  $R \in \mathbb{F}(X \times Y)$ ,  $S \in \mathbb{F}(Y \times Z)$  and  $\wedge_{\circ}$  some fuzzy conjunction. Then the  $\circ$ -composition („ $\circ$ -složená relace“) is

$$R \circ S(x, z) = \bigvee_{y \in Y}^S R(x, y) \wedge_{\circ} S(y, z) \quad (20)$$

1. For infinite domains,  $\bigvee^S$  is computed using the  $\sup$  instead of  $\max$ .
2. Instead of the “for some  $y$ ” in *crisp relations*, the disjunction “finds such a  $y$ ” that maximizes the conjunction.

## Example of a fuzzy relation

$$R(x, y) = \begin{cases} x + y & x, y \in [0, \frac{1}{2}] \\ 0 & \text{otherwise} \end{cases}$$

$$S(x, y) = \begin{cases} x \cdot y & x, y \in [0, 1] \\ 0 & \text{otherwise} \end{cases}$$

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

**property**

**using set axioms**

---



# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using set axioms
reflexive	$E \subseteq R$

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
$\circ$ -anti-symmetric	$R \cap R^{-1} \subseteq E$

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
$\circ$ -anti-symmetric	$R \cap_{\circ} R^{-1} \subseteq E$
$\circ$ -transitive	$R \circ_{\circ} R \subseteq R$

# Properties of fuzzy relations

Then the relation  $R \subseteq \Delta \times \Delta$  is called

property	using set axioms
reflexive	$E \subseteq R$
symmetric	$R = R^{-1}$
$\circ$ -anti-symmetric	$R \cap_{\circ} R^{-1} \subseteq E$
$\circ$ -transitive	$R \circ_{\circ} R \subseteq R$
$\circ$ -partial order	reflexive, $\circ$ -transitive and $\circ$ -anti-symmetric
$\circ$ -equivalence	reflexive, $\circ$ -transitive and $\circ$ -symmetric

# Properties in a finite domain

If the universe  $\Delta$  is finite, the relation can be written as a matrix. Their properties are reflected in the relation's matrix:

- **Reflexivity:** Cells on the main diagonal ?.
- **Symmetry:** Cells symmetric over the main diagonal ?.
- **Anti-symmetry:** Cells symmetric over the main diagonal ?.
  - For *S*- and *A*-anti-symmetry, ?.
  - For *L*-anti-symmetry, ?.
- **Transitivity:** More difficult (see example on the next slide).

# Properties in a finite domain

If the universe  $\Delta$  is finite, the relation can be written as a matrix. Their properties are reflected in the relation's matrix:

- **Reflexivity:** Cells on the main diagonal are 1.
- **Symmetry:** Cells symmetric over the main diagonal are equal.
- **Anti-symmetry:** Cells symmetric over the main diagonal have conjunction equal to zero.
  - For S- and A-anti-symmetry, one of the elements must be zero.
  - For L-anti-symmetry, their sum must be less or equal to 1.
- **Transitivity:** More difficult (see example on the next slide).

# Example on fuzzy relation properties

Let  $\Delta = \{A, B, C, D\}$  and  $R \in \mathbb{F}(\Delta \times \Delta)$ .

$R$	A	B	C	D
A		0.5		0.1
B			0.2	
C				
D		0.2		

Fill the missing cells in the table to make  $R$

- S-equivalence
- A-equivalence



## Theorem 264.

Let  $R$ ,  $S$  and  $T$  be relations (defined over sets that “make sense”) The following equations hold:

## Theorem 264.

Let  $R$ ,  $S$  and  $T$  be relations (defined over sets that “make sense”) The following equations hold:

$$R \circ E = R, \quad E \circ R = R \quad (21)$$

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1} \quad (22)$$

$$R \circ (S \circ T) = (R \circ S) \circ T \quad (23)$$

$$\left( R \bigcap^S S \right) \circ T = (R \circ T) \circ (S \circ T) \quad (24)$$

$$R \circ \left( S \bigcap^S T \right) = (R \circ S) \circ (R \circ T) \quad (25)$$

## Theorem 264.

Let  $R$ ,  $S$  and  $T$  be relations (defined over sets that “make sense”) The following equations hold:

$$R \circ E = R, \quad E \circ R = R \quad (21)$$

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1} \quad (22)$$

$$R \circ (S \circ T) = (R \circ S) \circ T \quad (23)$$

$$(R \bigcap^S S) \circ T = (R \circ T) \circ (S \circ T) \quad (24)$$

$$R \circ (S \bigcap^S T) = (R \circ S) \circ (R \circ T) \quad (25)$$

(21) describes the *identity element*, (22) the *inverse of composition*, (23) is the *asociativity*, (24) and (25) the *right-* and *left-distributivity*.

## Proof of 264.

Proving (21) and (22) is trivial.

$$"R \circ (S \circ T)"(x, w) = \bigvee_y^s R(x, y) \wedge "S \circ T"(y, w) \quad (26)$$

$$= \bigvee_y^s R(x, y) \wedge \left( \bigvee_z^s S(y, z) \wedge T(z, w) \right) \quad (27)$$

$$= \bigvee_y^s \bigvee_z^s R(x, y) \wedge S(y, z) \wedge T(z, w) \quad (28)$$

$$= \bigvee_z^s \bigvee_y^s R(x, y) \wedge S(y, z) \wedge T(z, w) \quad (29)$$

## Proof of 264 (contd.).

$$= \bigvee_z^s \bigvee_y^s R(x, y) \underset{\circ}{\wedge} S(y, z) \underset{\circ}{\wedge} T(z, w) \quad (30)$$

$$= \bigvee_z^s \left( \bigvee_y^s R(x, y) \underset{\circ}{\wedge} S(y, z) \right) \underset{\circ}{\wedge} T(z, w) \quad (31)$$

$$= \bigvee_z^s "R \underset{\circ}{S}"(x, z) \underset{\circ}{\wedge} T(z, w) \quad (32)$$

$$= "R \underset{\circ}{S} \underset{\circ}{T}"(x, w) \quad (33)$$

Proof of (24) and (25) is similar (uses the distributivity law), only shorter. See [Navara and Olšák, 2001] for details.

## Extensions: Sometimes it is useful to consider...

- ...a  $\varepsilon$ -*reflective* relation

$$R(\mathbf{x}, \mathbf{x}) \geq \varepsilon \quad (34)$$

## Extensions: Sometimes it is useful to consider...

- ...a  $\varepsilon$ -*reflective* relation

$$R(\mathbf{x}, \mathbf{x}) \geq \varepsilon \quad (34)$$

- ...a *weakly reflexive* relation

$$R(\mathbf{x}, \mathbf{y}) \leq R(\mathbf{x}, \mathbf{x}) \text{ and } R(\mathbf{y}, \mathbf{x}) \leq R(\mathbf{x}, \mathbf{x}) \text{ for all } \mathbf{x}, \mathbf{y} \quad (35)$$

## Extensions: Sometimes it is useful to consider...

- ...a  $\varepsilon$ -*reflective* relation

$$R(x, x) \geq \varepsilon \quad (34)$$

- ...a *weakly reflexive* relation

$$R(x, y) \leq R(x, x) \text{ and } R(y, x) \leq R(x, x) \text{ for all } x, y \quad (35)$$

- Relation is 1-reflective iff reflexive.
- If a relation is reflexive, then it is weakly reflexive.



# Extensions: Sometimes it is useful to consider...

- ...a *non-involutive negation* by refusing (N2)

$$\neg\neg\alpha \neq \alpha$$

and adopting a weaker axiom

$$\neg\neg 0 = 1 \text{ and } \neg\neg 1 = 0 \quad (\text{N0})$$

## Example

Gödel negation

$$\neg_G \alpha = \begin{cases} 1 & \alpha = 0 \\ 0 & \text{otherwise} \end{cases} \quad (36)$$



Navara, M. and Olšák, P. (2001).  
Základy fuzzy množin.  
Nakladatelství ČVUT.