

AE4M33RZN, Fuzzy logic: Tutorial examples

Radomír Černocho

radomir.cernoch@fel.cvut.cz

2015

Faculty of Electrical Engineering, CTU in Prague

Task 2

Assignment

On the universe $\Delta = \{a, b, c, d\}$ there is a fuzzy set

$$\mu_A = \{(a; 0.3), (b; 1), (c; 0.5)\}.$$

Find its horizontal representation.

$$R_A(\alpha) = \begin{cases} X & \alpha = 0, \\ \{a, b, c\} & \alpha \in (0; 0.3), \\ \{b, c\} & \alpha \in (0.3; 0.5), \\ \{b\} & \alpha \in (0.5; 1). \end{cases}$$

Assignment

The fuzzy set A has a horizontal representation

$$R_A(\alpha) = \begin{cases} \{a, b, c, d\} & \alpha \in \langle 0, 1/3 \rangle \\ \{a, d\} & \alpha \in (1/3, 1/2) \\ \{d\} & \alpha \in (1/2, 2/3) \\ \emptyset & \alpha \in (2/3, 1) \end{cases}$$

Find the vertical representation.

Solution

$$\mu_A(a) = \sup\{\alpha \in \langle 0, 1 \rangle : a \in R_A(\alpha)\} = \sup\langle 0, 1/2 \rangle = 1/2$$

$$\mu_A(b) = 1/3, \quad \mu_A(c) = 1/3, \quad \mu_A(d) = 2/3, \quad \text{therefore}$$

$$\mu_A = \{(a, 1/2), (b, 1/3), (c, 1/3), (d, 2/3)\}$$

Task 5

Assignment

On the universe $\Delta = \mathbb{R}$ there is a fuzzy set A :

$$\mu_A(x) = \begin{cases} x & x \in \langle 0; 1 \rangle \\ 2 - x & x \in (1; 1.5) \\ 0 & \text{otherwise} \end{cases}$$

Find its horizontal representation.

$$R_A(\alpha) = \begin{cases} \mathbb{R} & \alpha = 0 \\ \langle \alpha; 1.5 \rangle & \alpha \in (0; 0.5) \\ \langle \alpha; 2 - \alpha \rangle & \alpha \in (0.5; 1) \end{cases}$$

Assignment

The fuzzy set A has a horizontal representation

$$R_A(\alpha) = \begin{cases} \mathbb{R} & \alpha = 0 \\ \langle \alpha^2, 1 \rangle & \text{otherwise.} \end{cases}$$

Find the vertical representation.

$$\mu_A(x) = \begin{cases} \sqrt{x} & x \in (0, 1) \\ 0 & \text{otherwise.} \end{cases}$$

Assignment

Decide if the following function is a fuzzy conjunction.

$$\alpha \underset{\circ}{\wedge} \beta = \begin{cases} \alpha & \beta = 1, \\ \beta & \alpha = 1, \\ \alpha\beta & \alpha\beta \geq 1/10, \max(\alpha, \beta) < 1, \\ 0 & \text{otherwise.} \end{cases}$$

Solution

The first two possibilities ensure the boundary condition. Comutativity and monotonicity are trivially satisfied. If one of the arguments is 1, the associativity as well. For $\alpha, \beta, \gamma < 1$ the associativity follows from

$$\alpha \underset{\circ}{\wedge} (\beta \underset{\circ}{\wedge} \gamma) = \begin{cases} \alpha\beta\gamma & \alpha\beta\gamma \geq 1/10. \\ 0 & \text{otherwise,} \end{cases}$$

We get the same for $(\alpha \underset{\circ}{\wedge} \beta) \underset{\circ}{\wedge} \gamma$.

It is always a fuzzy conjunction (interpretable as an algebraic conjunction, in which we ignore small values, e.g. for filtering small values).

Assignment

Prove that the Sugeno class of negations are all fuzzy negations

$$\neg_{s_\lambda} \alpha = \frac{1 - \alpha}{1 + \lambda \alpha} \text{ for } \lambda \in (-1, \infty,).$$

Solution

When canceling out terms in a fraction, we use $1 + \lambda\alpha > 0$.

- **Involutivity:** Assuming $\lambda > -1$:

$$\overline{s_\lambda} \overline{s_\lambda} \alpha = \frac{1 - \frac{1-\alpha}{1+\lambda\alpha}}{1 + \lambda \frac{1-\alpha}{1+\lambda\alpha}} = \frac{1 + \lambda\alpha - 1 + \alpha}{1 + \lambda\alpha + \lambda - \lambda\alpha} = \frac{\alpha(\lambda + 1)}{\lambda + 1} = \alpha$$

- **Non-increasing:** If $\alpha \leq \beta$, then

$$\frac{1 - \alpha}{1 + \lambda\alpha} \geq \frac{1 - \beta}{1 + \lambda\beta}$$

$$(1 + \lambda\beta)(1 - \alpha) \geq (1 + \lambda\alpha)(1 - \beta)$$

$$\beta \geq \alpha$$

Task 12

Assignment

Decide if the following function is a fuzzy conjunction.

$$\alpha \diamond \beta = \begin{cases} \alpha\beta & \alpha\beta \geq 0,01 \text{ or } \max(\alpha, \beta) = 1, \\ 0 & \text{otherwise} \end{cases}$$

Assignment

Decide if the following function is a fuzzy conjunction.

$$\underset{\circ}{\wedge}: \langle 0, 1 \rangle^2 \rightarrow \langle 0, 1 \rangle$$

$$\alpha \underset{\circ}{\wedge} \beta = \begin{cases} \min(\alpha, \beta) & \alpha + \beta \geq 1 \\ 0 & \text{otherwise} \end{cases}$$

Assignment

Decide, if for all $\alpha, \beta \in [0, 1]$ holds $(\alpha \overset{\circ}{\vee} \beta) \overset{\circ}{\wedge} (\alpha \overset{\circ}{\vee} \neg \beta) = \alpha$, where the

disjunction $\overset{\circ}{\vee}$ is

1. standard, $\overset{S}{\vee}$,
2. algebraic, $\overset{\Delta}{\vee}$.
3. Łukasiewicz, $\overset{L}{\vee}$,

Solution

1. No. Counterexample: $\alpha = 0.5, \beta = 0.1$.
2. Yes: $\max(0, (\alpha + \beta - \alpha\beta) + (\alpha + (1 - \beta) - \alpha(1 - \beta)) - 1) =$
 $= \max(0, \alpha) = \alpha$
3. No. Counterexample: $\alpha = 0.9, \beta = 0.8$.

Assignment

Decide if the function $\underset{\circ}{\wedge} : \langle 0, 1 \rangle^2 \rightarrow \langle 0, 1 \rangle$

$$\alpha \underset{\circ}{\wedge} \beta = \begin{cases} \alpha \beta & \alpha + \beta \geq 1 \\ 0 & \text{otherwise} \end{cases}$$

is a fuzzy conjunction.

Assignment

Decide if the $(\alpha \wedge \alpha) \vee (\alpha \wedge \alpha) \leq \alpha$ holds for

1. standard, \mathbb{S}
2. algebraic, \mathbb{A}
3. Łukasiewicz, \mathbb{L}

Assignment

Decide, which equalities hold:

$$1. (\alpha \underset{S}{\wedge} \alpha) \underset{L}{\vee} (\alpha \underset{S}{\wedge} \beta) = \alpha \underset{S}{\wedge} (\alpha \underset{L}{\vee} \beta)$$

$$2. (\alpha \underset{L}{\wedge} \alpha) \underset{S}{\vee} (\alpha \underset{L}{\wedge} \beta) = \alpha \underset{L}{\wedge} (\alpha \underset{S}{\vee} \beta)$$

$$3. \alpha \underset{S}{\vee} (\alpha \underset{L}{\wedge} \beta) = \alpha \underset{L}{\wedge} (\alpha \underset{S}{\vee} \beta)$$

Justify your conclusions.

Assignment

Decide, which equalities hold:

$$1. (\alpha \underset{S}{\wedge} \beta) \underset{L}{\wedge} \gamma = \alpha \underset{S}{\wedge} (\beta \underset{L}{\wedge} \gamma)$$

$$2. \neg \underset{S}{\wedge} (\alpha \underset{A}{\vee} \beta) = \neg \underset{S}{\wedge} \alpha \underset{L}{\wedge} \neg \underset{S}{\wedge} \beta$$

$$3. (\alpha \underset{L}{\wedge} \alpha) \underset{V}{\vee} \neg \underset{S}{\wedge} \alpha = (\neg \underset{S}{\wedge} \alpha \underset{L}{\wedge} \neg \underset{S}{\wedge} \alpha) \underset{V}{\vee} \alpha$$

Justify your conclusions.

Assignment

Verify that $\alpha \underset{\circ}{\wedge} (\alpha \underset{\circ}{\overset{R}{\Rightarrow}} \beta) = \alpha \underset{S}{\wedge} \beta$ holds for

1. algebraic ops.
2. standard ops.

We will show the solution for algebraic operations. The other ones are similar.

$$\alpha \underset{A}{\wedge} (\alpha \underset{A}{\overset{R}{\rightarrow}} \beta) = \left\{ \begin{array}{l} \alpha \cdot 1 = \alpha \text{ for } \alpha \leq \beta \\ \alpha \cdot \frac{\beta}{\alpha} = \beta \text{ for } \alpha > \beta \end{array} \right\} = \min(\alpha, \beta) = \alpha \underset{S}{\wedge} \beta$$

Task 22

Assignment

Complete the table, so that R is a S -partial order.

R	a	b	c	d
a				
b	0.5			
c		0.3		
d		0.2		

Solution

Reflexivity implies 1's on the diagonal. S-partial order implies 0's to non-zero elements symmetric over the main diagonal.

R	a	b	c	d
a	1	0	x'	y'
b	0.5	1	0	0
c	x	0.3	1	z'
d	y	0.2	z	1

The transitivity implies e.g. $R(3, 2) \underset{S}{\wedge} R(2, 1) \leq R(3, 1)$, which translates

into a condition $\min(0.3, 0.5) \leq x$

Using this and similar conditions, we derive the subspace of all solutions: $z \leq 0.2$, $x \geq 0.3$, $y \geq 0.2$, $\min(y, z') \leq a$, $x' = y' = 0$.

Ex: Jim revisited

We will use the Lukasiewicz logic in the following examples ($\sqcap = \sqcap, \dots$).

$$\langle \mathit{jim} : \mathit{Male} \mid 0.9 \rangle \quad (1)$$

$$\langle \mathit{jim} : \mathit{Female} \mid 0.2 \rangle \quad (2)$$

$$\langle \mathit{Male} \sqcap \mathit{Female} \sqsubseteq \perp \mid 1 \rangle \quad (3)$$

Ex: Jim revisited

We will use the Lukasiewicz logic in the following examples ($\sqcap = \sqcap, \dots$).

$$\langle jim : \text{Male} \mid 0.9 \rangle \quad (1)$$

$$\langle jim : \text{Female} \mid 0.2 \rangle \quad (2)$$

$$\langle \text{Male} \sqcap \text{Female} \sqsubseteq \perp \mid 1 \rangle \quad (3)$$

The interpretation domain is $\Delta^{\mathcal{I}_1} = \Delta^{\mathcal{I}_2} = \{j\}$, $jim^{\mathcal{I}_1} = jim^{\mathcal{I}_2} = j$.

$$\text{Male}^{\mathcal{I}_1} = \{(j; 0.9)\}$$

$$\text{Male}^{\mathcal{I}_2} = \{(j; 0.9)\}$$

$$\text{Female}^{\mathcal{I}_1} = \{(j; 0)\}$$

$$\text{Female}^{\mathcal{I}_2} = \{(j; 0.2)\}$$

Ex: Jim revisited (check your knowledge)

Let's check the interpretation against the definitions...

$\mathcal{I} \models \tau$	$\tau_{(1)}$	$\tau_{(2)}$	$\tau_{(3)}$
\mathcal{I}_1	?	?	?
\mathcal{I}_2	?	?	?

Ex: Jim revisited (check your knowledge)

Let's check the interpretation against the definitions...

$\mathcal{I} \models \tau$	$\tau_{(1)}$	$\tau_{(2)}$	$\tau_{(3)}$
\mathcal{I}_1	yes	no	yes
\mathcal{I}_2	yse	yes	no

Ex: Jim revisited (in fuzzyDL)

Let's change the weights and encode the example in fuzzyDL:

```
(instance jim Male 0.4)
```

```
(instance jim Female 0.2)
```

```
(1-implies (and Male Female) *bottom* 0.9)
```

```
(min-instance? jim Male)
```

```
(max-instance? jim Male)
```

```
(min-instance? jim Female)
```

```
(max-instance? jim Female)
```

Let $\langle jim : \text{Male} | \alpha \rangle$ and $\langle jim : \text{Female} | \beta \rangle$, what are the bounds on α and β ? fuzzyDL shows that $0.4 \leq \alpha \leq 0.9$ and $0.2 \leq \beta \leq 0.7$. Why?

Ex: Smokers

Recall the motivational example from the first lecture:

$$\langle \text{symmetric}(\text{friend}) \rangle \quad (4)$$

$$\langle (\text{anna}, \text{bill}) : \text{friend} \mid 1 \rangle \quad (5)$$

$$\langle (\text{bill}, \text{cloe}) : \text{friend} \mid 1 \rangle \quad (6)$$

$$\langle (\text{cloe}, \text{dirk}) : \text{friend} \mid 1 \rangle \quad (7)$$

$$\langle \text{anna} : \text{Smoker} \mid 1 \rangle \quad (8)$$

$$\langle \exists \text{friend} \cdot \text{Smoker} \sqsubseteq \text{Smoker} \mid 0.7 \rangle \quad (9)$$

What are the bounds on $\langle i : \text{Smoker} \rangle$ for $i \in \{\text{anna}, \text{bill}, \text{cloe}, \text{dirk}\}$?

Ex: Smokers

What changes if we add

$$\langle \text{dirk} : \neg \text{Smoker} \mid 0.7 \rangle \quad (10)$$

(11)

What are the bounds on $\langle i : \neg \text{Smoker} \rangle$ for $i \in \{\text{anna}, \text{bill}, \text{cloe}, \text{dirk}\}$?

Ex: Smokers (in fuzzyDL)

```
(implies (some friendOf Smoker) Smoker 0.7)
```

```
(symmetric friendOf)  
(related anna bill friendOf)  
(related bill cloe friendOf)  
(related cloe dirk friendOf)
```

```
(instance anna Smoker)  
(instance dirk (not Smoker) 0.7)
```

```
(min-instance? anna Smoker)  
(min-instance? bill Smoker)  
(min-instance? cloe Smoker)  
(min-instance? dirk Smoker)
```

```
(max-instance? anna Smoker)  
(max-instance? bill Smoker)  
(max-instance? cloe Smoker)  
(max-instance? dirk Smoker)
```

Concrete data types

The domain $\Delta^{\mathcal{F}}$ is an unordered set. This is good for modelling categorical data: e.g. colors, people, ...

General idea: Extended interpretation

But we also need to include real numbers \mathbb{R} . The *fuzzy description logic with concrete datatypes* $\mathcal{R}\mathcal{G}\mathcal{H}\mathcal{E}$ —uses “abstract objects” and “concrete objects”:

$$\Delta^{\mathcal{F}} = \Delta_a^{\mathcal{F}} \cup \mathbb{R}$$

Concrete data types

- *Concrete individuals*, are interpreted as objects from \mathbb{R} .

Concrete data types

- *Concrete individuals*, are interpreted as objects from \mathbb{R} .
- *Concrete concepts*, are interpreted as subsets from \mathbb{R} .

Concrete data types

- *Concrete individuals*, are interpreted as objects from \mathbb{R} .
- *Concrete concepts*, are interpreted as subsets from \mathbb{R} .
- *Concrete roles*, are interpreted as subsets from $(\Delta_a^{\mathcal{F}} \times \mathbb{R})$.

Concrete data types

- *Concrete individuals*, are interpreted as objects from \mathbb{R} .
- *Concrete concepts*, are interpreted as subsets from \mathbb{R} .
- *Concrete roles*, are interpreted as subsets from $(\Delta_a^{\mathcal{F}} \times \mathbb{R})$.

All non-concrete notions are called *abstract*.

Concrete data types: New concepts




Fig. 1. (a) Trapezoidal function; (b) Triangular function; (c) *L*-function; (d) *R*-function; (e) Crisp interval; (f) Linear function.

Ex: Age of parents

```
(related adam bob parent) (related adam eve parent)

(define-fuzzy-concept around23 triangular(0,100, 18,23,26))
(define-fuzzy-concept moreTh17 right-shoulder(0,100, 13,21))
(instance bob (some age around23) 0.9)
(instance eve (some age moreTh17))

(define-fuzzy-concept young left-shoulder(0,100, 17,25))
(define-concept YoungPerson (some age young))

(min-instance? eve YoungPerson) (max-instance? eve YoungPerson)
(min-instance? bob YoungPerson) (max-instance? bob YoungPerson)
(min-instance? adam (all parent YoungPerson))
(max-instance? adam (all parent YoungPerson))
(min-instance? adam (some parent YoungPerson))
(max-instance? adam (some parent YoungPerson))
```

Ex: Age of parents

1. What are the bounds on α from $\langle \text{eve} : \text{YoungPerson} \mid \alpha \rangle$?

Ex: Age of parents

1. What are the bounds on α from $\langle \text{eve} : \text{YoungPerson} \mid \alpha \rangle$?

Start by drawing the concept **around23**, then construct an interpretation. How much freedom do you have when constructing the interpretation?

Ex: Age of parents

1. What are the bounds on α from $\langle \text{eve} : \text{YoungPerson} \mid \alpha \rangle$?

Start by drawing the concept `around23`, then construct an interpretation. How much freedom do you have when constructing the interpretation?

2. Let fuzzyDL reasoner give you both bounds on $\langle i : \text{YoungPerson} \mid \beta_i \rangle$ for $i \in \{\text{eve}, \text{bob}\}$.

How do you infer the bounds on $\langle \text{adam} : \text{YoungPerson} \mid \gamma \rangle$?

Ex: Car dealing

1. The buyer wants a **passenger** that costs **less than €26000**.
2. If there is an **alarm system** in the car, **then** he is satisfied with paying no more than **€22300**, but he can go up to **€22750** with a lesser degree of satisfaction.
3. The **driver insurance**, **air conditioning** and the **black color** are important factors.
4. Preferably the price is no more than **€22000**, but he can go to **€24000** to a lesser degree of satisfaction.

Ex: Car dealing

1. The seller wants to sell no less than **€22000**.
2. Preferably the buyer buys the **insurance plus** package.
3. If the **color is black**, then it is highly possible the car has an **air-conditioning**.

This can be formalized in fuzzy description logic.

Ex: Car dealing

1. The seller wants to sell no less than **€22000**.
2. Preferably the buyer buys the **insurance plus** package.
3. If the **color is black**, then it is highly possible the car has an **air-conditioning**.

This can be formalized in fuzzy description logic.

We have the background knowledge:

$\langle \text{Sedan} \sqsubseteq \text{PassengerCar} \mid 1 \rangle$

$\langle \text{InsurancePlus} = \text{DriverInsurance} \sqcap \text{TheftInsurance} \mid 1 \rangle$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$
3. $B_2 = \text{DriverInsurance},$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$
3. $B_2 = \text{DriverInsurance}, B_3 = \text{AirCondition},$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$
3. $B_2 = \text{DriverInsurance}, B_3 = \text{AirCondition}, B_4 = \exists \text{color} \cdot \text{Black}$
4. $B_5 = \exists \text{price} \cdot \text{I.sh.}(22000, 24000)$

The buyer's preferences:

1. $S = \text{PassengerCar} \sqcap \exists \text{price} \cdot \geq 22000$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$
3. $B_2 = \text{DriverInsurance}, B_3 = \text{AirCondition}, B_4 = \exists \text{color} \cdot \text{Black}$
4. $B_5 = \exists \text{price} \cdot \text{I.sh.}(22000, 24000)$

The buyer's preferences:

1. $S = \text{PassengerCar} \sqcap \exists \text{price} \cdot \geq 22000$
2. $S_1 = \text{InsurancePlus}$

Ex: Car dealing

The buyer's preferences:

1. $B = \text{PassengerCar} \sqcap \exists \text{price} \cdot \leq 26000$
2. $B_1 = \text{AlarmSystem} \mapsto \exists \text{price} \cdot \text{I.sh.}(22300, 22750)$
3. $B_2 = \text{DriverInsurance}, B_3 = \text{AirCondition}, B_4 = \exists \text{color} \cdot \text{Black}$
4. $B_5 = \exists \text{price} \cdot \text{I.sh.}(22000, 24000)$

The buyer's preferences:

1. $S = \text{PassengerCar} \sqcap \exists \text{price} \cdot \geq 22000$
2. $S_1 = \text{InsurancePlus}$
3. $S_2 = (0.5 (\exists \text{color} \cdot \text{Black}) \mapsto \text{AirCondition})$

Ex: Car dealing

We know that S and B are hard constraints and $B_{1..5}$ and $S_{1..2}$ are soft preferences. All the concepts can be “summed up”:

Ex: Car dealing

We know that S and B are hard constraints and $B_{1..5}$ and $S_{1..2}$ are soft preferences. All the concepts can be “summed up”:

$$\text{Buy} = B \sqcap (0.1B_1 + 0.2B_2 + 0.1B_3 + 0.4B_4 + 0.2B_5)$$

and

$$\text{Sell} = S \sqcap (0.6S_1 + 0.4S_2)$$

Ex: Car dealing

We know that S and B are hard constraints and $B_{1..5}$ and $S_{1..2}$ are soft preferences. All the concepts can be “summed up”:

$$\text{Buy} = B \sqcap (0.1B_1 + 0.2B_2 + 0.1B_3 + 0.4B_4 + 0.2B_5)$$

and

$$\text{Sell} = S \sqcap (0.6S_1 + 0.4S_2)$$

A good choice of \sqcap can make B a hard constraint.

Ex: Car dealing

Optimal match

$$bsb(K, \text{Buy} \sqcap \text{Sell})$$

Finds the optimal match between a seller and a buyer. (Finds an ideal, imaginary car that maximizes satisfaction of both parties.)

Particular car

$$bdb(K, \langle \text{audiTT} : \text{Buy} \sqcap \text{Sell} \rangle)$$

Finds the degree of satisfaction for a particular car *audiTT*.