

GRAPHICAL MARKOV MODELS (WS2023)
2. SEMINAR

Assignment 1 (*Galton-Watson-Process – a population model*). Individuals of a certain population can have $n = 0, 1, 2, \dots$ offspring at the end of their life. The corresponding probabilities are c_0, c_1, c_2, \dots . Let s_i denote the size of the population in the i -th generation.

a) Model the process as a Markov chain. Deduce a formula for the transition probabilities $p(s_i = k \mid s_{i-1} = m)$.

b)** Calculate the extinction probability ρ_k , i.e. the probability that the population will eventually extinct if it starts with k individuals in the first generation.

Hints:

- (1) Express ρ_k in terms of $\rho := \rho_1$.
- (2) Try to find a functional relationship for ρ and the probabilities $c_k, k = 0, 1, 2, \dots$
- (3) Analyse the resulting fix-point equation for ρ .

Let us consider the following standard Markov chain model for the next two assignments. The probability for sequences $s = (s_1, \dots, s_n)$ of length n with states $s_i \in K$ is given by:

$$p(s) = p(s_1) \prod_{i=2}^n p(s_i \mid s_{i-1}).$$

The conditional probabilities $p(s_i \mid s_{i-1})$ and the marginal probability $p(s_1)$ for the first element are assumed to be known.

Assignment 2. Suppose that there is a special state $k^* \in K$. We want to know how often this state appears on average in a sequence generated by the model. Describe an efficient method for computing this average.

Hint: Use the fact that the expected value of a sum of random variables is equal to the sum of their expected values.

Assignment 3. Let $A \subset K$ be a subset of states and let $\mathcal{A} = A^n$ denote the set of all sequences s with $s_i \in A$ for all $i = 1, \dots, n$. Find an efficient algorithm for computing the probability $p(\mathcal{A})$ of the event \mathcal{A} .

Assignment 4. According to Definition 1b of Sec. 1 of the lecture, any Markov chain model can be specified in the form

$$p(s) = \frac{1}{Z} \prod_{i=2}^n g_i(s_{i-1}, s_i)$$

with arbitrary functions $g_i: K^2 \rightarrow \mathbb{R}_+$ and the normalisation constant Z . Find an algorithm for computing the pairwise marginal probabilities $p(s_{i-1} = k, s_i = k')$ for all $k, k' \in K$ and all $i = 2, \dots, n$ from the given functions $g_i, i = 2, \dots, n$.

Assignment 5*: Suppose that a regular language \mathcal{L} of strings over the finite alphabet Σ is described by a non-deterministic finite-state machine. Given a string $y \notin \mathcal{L}$, the task is to find the string $x \in \mathcal{L}$ with smallest Hamming distance to y , i.e.

$$x^* = \arg \min_{x \in \mathcal{L}} d_h(x, y),$$

where d_h denotes the Hamming distance. Construct an efficient algorithm for solving this task.